

FLEA HEAD, NERVE CORD, HINDGUT AND MALPIGHIAN TUBULE  
NUCLEIC ACID MOLECULES, PROTEINS AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

5           This application claims priority to U.S. Provisional Patent Application Serial  
No. 60/319,414, filed July 22, 2002, entitled "FLEA HEAD, NERVE CORD, HINDGUT  
AND MALPIGHIAN TUBULE NUCLEIC ACID MOLECULES, PROTEINS AND  
USES THEREOF," which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

10           The present invention relates to nucleic acid molecules isolated from the head and  
nerve cord of a flea, nucleic acid molecules isolated from the hindgut and Malpighian  
tubule of a flea, proteins encoded by such nucleic acid molecules, antibodies raised  
against such proteins, and inhibitors of such proteins. The present invention also includes  
therapeutic compositions comprising such nucleic acid molecules, proteins, antibodies,  
15   and/or other inhibitors, as well as uses thereof.

BACKGROUND OF THE INVENTION

Flea infestation of animals is a health and economic concern to pet owners. In  
particular, the bites of fleas are a problem for animals maintained as pets because the  
infestation becomes a source of annoyance not only for the pet but also for the pet owner  
20   who may find his or her home generally contaminated with insects. Fleas are known to  
directly cause a variety of diseases, including allergies, and also carry a variety of  
infectious agents including, but not limited to, endoparasites (e.g., nematodes, cestodes,  
trematodes and protozoa), bacteria and viruses. As such, fleas are a problem not only  
when they are on an animal but also when they are in the general environment of the  
25   animal.

Bites from fleas are a particular problem on many animals because they can cause  
a hypersensitive response in animals which is manifested as disease. For example, bites  
from fleas can cause an allergic disease called flea allergic (or allergy) dermatitis (FAD).

A hypersensitive response in animals typically results in localized tissue inflammation and damage, causing substantial discomfort to the animal.

The medical importance of flea infestation has prompted the development of reagents capable of controlling flea infestation. Commonly encountered methods to control flea infestation are generally focused on use of insecticides, which are often  
5 unsuccessful for one or more of the following reasons: (1) failure of owner compliance (frequent administration is required); (2) behavioral or physiological intolerance of the pet to the pesticide product or means of administration; and (3) the emergence of flea populations resistant to the prescribed dose of pesticide.

10 Thus, there remains a need to develop a reagent and a method to protect animals from flea infestation.

#### SUMMARY OF THE INVENTION

The present invention relates to a novel product and process for protection of animals from flea infestation.

15 The present invention provides flea head and nerve cord (HNC) proteins and flea hindgut and Malpighian tubule (HMT) proteins; nucleic acid molecules encoding flea HNC proteins and flea HMT proteins; antibodies raised against such proteins (i.e., anti-flea HNC antibodies and anti-flea HMT antibodies respectively); mimetopes of such proteins or antibodies; and compounds that inhibit flea HNC or HMT activity (i.e.,  
20 inhibitory compounds or inhibitors).

The present invention also includes methods to obtain such proteins, mimetopes, nucleic acid molecules, antibodies and inhibitory compounds. The present invention also includes the use of proteins and antibodies to identify such inhibitory compounds as well as assay kits to identify such inhibitory compounds. Also included in the present  
25 invention are therapeutic compositions comprising proteins, mimetopes, nucleic acid molecules, antibodies and inhibitory compounds of the present invention including protective compounds derived from a protein of the present invention that inhibit the

activity of HNC and/or HMT proteins; also included are uses of such therapeutic compounds to reduce flea infestation.

One embodiment of the present invention is an isolated nucleic acid molecule that hybridizes with a nucleic acid molecule selected from the group consisting of a nucleic acid sequence of Table I and/or Table II, or a nucleic acid sequence complementary to a  
5 nucleic acid sequence of Table I and/or Table II under conditions that allow less than or equal to 30% base pair mismatch.

Another embodiment of the present invention is an isolated nucleic acid molecule having a nucleic acid sequence that is at least 70% identical to a nucleic acid sequence of  
10 Table I and/or Table II or complements thereof as well as fragments of such sequences.

The present invention also relates to recombinant molecules, recombinant viruses and recombinant cells that include a nucleic acid molecule of the present invention. Also included are methods to produce such nucleic acid molecules, recombinant molecules, recombinant viruses and recombinant cells.

15 Another embodiment of the present invention includes an isolated flea HMT and/or HNC protein that is at least 70% identical to an amino acid sequence encoded by a nucleic acid sequence of Table I and/or Table II, and fragments thereof, wherein such fragments can elicit an immune response against respective flea proteins or have activity comparable to respective flea proteins.

20 Another embodiment of the present invention includes an isolated protein encoded by a nucleic acid molecule that hybridizes with the complement of a nucleic acid sequence of Table I and/or Table II, under conditions that allow less than or equal to 30% base pair mismatch.

#### DETAILED DESCRIPTION OF THE INVENTION

25 The present invention provides for nucleic acid molecules isolated from the head and/or nerve cord of a flea, nucleic acid molecules isolated from the hindgut and/or Malpighian tubule of a flea, proteins encoded by such nucleic acid molecules, antibodies

raised against such proteins, and inhibitors of such proteins. As used herein, nucleic acid molecules isolated from the head and/or nerve cord of a flea and proteins encoded by such nucleic acid molecules are also referred to as flea HNC, or HNC, nucleic acid molecules and proteins respectively; and nucleic acid molecules isolated from the hindgut and/or Malpighian tubules of a flea and proteins encoded by such nucleic acid molecules are referred to as flea HMT or HMT, nucleic acid molecules and proteins respectively. HNC nucleic acid molecules and HMT nucleic acid molecules of the present invention are nucleic acid molecules that are primarily expressed in flea HNC tissues and HMT tissues respectively, but which may be expressed in cells derived from flea tissues other than HNC and HMT. HNC and HMT nucleic acid molecules and proteins of the present invention can be isolated from a flea or prepared recombinantly or synthetically. HMT and HNC nucleic acid molecules of the present invention can be RNA or DNA; examples of nucleic acid molecules include, but are not limited to, complementary DNA (cDNA) molecules, genomic DNA molecules, synthetic DNA molecules, DNA molecules which are specific tags for messenger RNA derived from HMT and HNC tissues, and corresponding mRNA molecules. As such, a flea nucleic acid molecule of the present invention is not intended refer to an entire chromosome within which such a nucleic acid molecule is contained, however, a flea HMT or HNC cDNA of the present invention may include all regions such as regulatory regions that control production of flea peritrophin proteins encoded by such a cDNA (such as, but not limited to, transcription, translation or post-translation control regions) as well as the coding region itself, and any introns or non-translated coding regions. As used herein, the phrases "HMT and/or HNC protein" and "HMT and HNC protein" refer to a protein expressed by a flea HMT tissue, by a flea HNC tissue, or by both flea HMT and HNC tissues. As used herein, the phrases "HMT and/or HNC nucleic acid molecule" and "HMT and HNC nucleic acid molecule" refer to a nucleic acid molecule that can be isolated from a HMT cDNA library, from a HNC cDNA library, or from both libraries, or a gene corresponding thereto.

The present invention also provides for HMT and HNC DNA molecules that are specific tags for messenger RNA molecules derived from HMT and HNC tissues. Such DNA molecules can correspond to an entire or partial sequence of a messenger RNA, and therefore, a DNA molecule corresponding to such a messenger RNA molecule (i.e. a cDNA molecule), can encode a full-length or partial-length protein. A nucleic acid molecule encoding a partial-length protein can be used directly as a probe or indirectly to generate primers to identify and/or isolate a cDNA nucleic acid molecule encoding a corresponding, or structurally related, full-length protein. Such a partial cDNA nucleic acid molecule can also be used in a similar manner to identify a genomic nucleic acid molecule, such as a nucleic acid molecule that contains the complete gene including regulatory regions, exons and introns. Methods for using partial HMT and HNC cDNA molecules and sequences to isolate full-length transcripts and corresponding cDNA molecules are described in the examples herein below.

The proteins and nucleic acid molecules of the present invention can be obtained from their natural source, or can be produced using, for example, recombinant nucleic acid technology or chemical synthesis. Also included in the present invention is the use of these proteins and nucleic acid molecules as well as antibodies and inhibitory compounds thereto as therapeutic compositions to protect animals from flea infestation as well as in other applications, such as those disclosed below.

Flea HMT and HNC proteins and nucleic acid molecules of the present invention have utility because they represent novel targets for anti-arthropod vaccines and chemotherapeutic drugs. The products and processes of the present invention are advantageous because they enable the inhibition of arthropod development, metamorphosis, feeding, digestion and/or reproduction processes that involve HMT and/or HNC proteins.

The head and nerve cord of the flea, including antennae, brain, corpora cardiacum, corpora allata, and subesophageal and abdominal ganglion tissues are of

interest as such tissues are highly enriched for transcripts that encode neuronal and endocrine targets, as well as targets involved in chemosensory and mechanosensory reception. By sequencing cDNA fragments from a library enriched in flea head and nerve cord nucleic acid sequences (referred to herein as HNC nucleic acid sequences),  
5 genes, and their respective full-length coding regions, integrally involved with flea neuronal and endocrine function are identified. Once identified, these genes can be further characterized and specific interference strategies are designed. As such, flea HNC proteins and nucleic acid molecules of the present invention have utility because they represent novel targets for anti-arthropod vaccines and chemotherapeutic drugs.

10 Blood-feeding insects such as fleas ingest large quantities of blood relative to their body weight and, as such, are adapted to reduce the volume of the ingested blood meal through the rapid elimination of water. In addition, the concentrations of sodium, potassium, and chloride ions in the blood meal are greater than in the hemolymph of fleas, necessitating the excretion of excessive amounts of these ions. The active transport  
15 of these ions from the hemolymph into the lumens of the Malpighian tubules and the hindgut drives the passive transport of water and other hemolymph contents into these organs as well. While passing through these organs, waste products from the hemolymph are excreted and needed nutrients, water, and salts are reabsorbed. As such, interfering with these essential processes is an important strategy for developing a product for  
20 controlling flea populations. By sequencing cDNA fragments from a library enriched in hindgut and Malpighian tubule nucleic acid sequences (referred to herein as HMT nucleic acid sequences), genes integrally involved with these processes, and their respective full-length coding regions, are identified. Once identified, these genes are further characterized and specific interference strategies can be designed. As such, flea HMT  
25 proteins and nucleic acid molecules of the present invention have utility because they represent novel targets for anti-arthropod vaccines and chemotherapeutic drugs.

One embodiment of the present invention is an isolated protein that includes a flea HMT and/or HNC protein. It is to be noted that the term "a" or "an" entity refers to one or more of that entity; for example, a protein, a nucleic acid molecule, an antibody and a therapeutic composition refers to "one or more" or "at least one" protein, nucleic acid molecule, antibody and therapeutic composition respectively. As such, the terms "a" (or "an"), "one or more" and "at least one" can be used interchangeably herein. It is also to be noted that the terms "comprising", "including", and "having" can be used interchangeably. According to the present invention, an isolated, or biologically pure, protein, is a protein that has been removed from its natural milieu. As such, "isolated" and "biologically pure" do not necessarily reflect the extent to which the protein has been purified. An isolated protein of the present invention can be obtained from its natural source, can be produced using recombinant DNA technology, or can be produced by chemical synthesis.

As used herein, isolated flea HMT and/or HNC proteins of the present invention can be full-length proteins or any homologue of such proteins. An isolated protein of the present invention, including a homologue, can be identified in a straight-forward manner by the protein's ability to elicit an immune response against a flea HMT and/or HNC protein or by the protein's HMT and/or HNC activity. Examples of flea HMT and HNC homologue proteins include flea HMT and HNC proteins in which amino acids have been deleted (e.g., a truncated version of the protein, such as a peptide), inserted, inverted, substituted and/or derivatized (e.g., by glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitoylation, amidation and/or addition of glycerophosphatidyl inositol) such that the homologue includes at least one epitope capable of eliciting an immune response against a flea HMT or HNC protein, and/or of binding to an antibody directed against a flea HMT or HNC protein. That is, when the homologue is administered to an animal as an immunogen, using techniques known to those skilled in the art, the animal will produce an immune response against at least one

epitope of a natural flea HMT or HNC protein. The ability of a protein to effect an immune response can be measured using techniques known to those skilled in the art. As used herein, the term "epitope" refers to the smallest portion of a protein or other antigen capable of selectively binding to the antigen binding site of an antibody or a T cell receptor. It is well accepted by those skilled in the art that the minimal size of a protein epitope is about four to six amino acids. As is appreciated by those skilled in the art, an epitope can include amino acids that naturally are contiguous to each other as well as amino acids that, due to the tertiary structure of the natural protein, are in sufficiently close proximity to form an epitope. According to the present invention, an epitope includes a portion of a protein comprising at least 4 amino acids, at least 5 amino acids, at least 6 amino acids, at least 10 amino acids, at least 15 amino acids, at least 20 amino acids, at least 25 amino acids, at least 30 amino acids, at least 35 amino acids, at least 40 amino acids or at least 50 amino acids in length.

In one embodiment of the present invention a flea homologue protein has HMT or HNC activity, i.e. the homologue exhibits an activity similar to its natural counterpart. Methods to detect and measure such activities are known to those skilled in the art.

Flea HMT and/or HNC homologue proteins can be the result of natural allelic variation or natural mutation. Flea HMT and/or HNC protein homologues of the present invention can also be produced using techniques known in the art including, but not limited to, direct modifications to the protein or modifications to the gene encoding the protein using, for example, classic or recombinant DNA techniques to effect random or targeted mutagenesis.

Flea HMT and HNC proteins of the present invention are encoded by flea HMT and HNC nucleic acid molecules, respectively. As used herein, flea HMT and HNC nucleic acid molecules include nucleic acid sequences related to natural flea HMT and HNC genes, and, preferably, to *Ctenocephalides felis* HMT and HNC genes. As used herein, flea HMT and HNC genes include all regions such as regulatory regions that



control production of flea HMT and HNC proteins encoded by such genes (such as, but not limited to, transcription, translation or post-translation control regions) as well as the coding region itself, and any introns or non-translated coding regions. As used herein, a nucleic acid molecule that “includes” or “comprises” a sequence may include that

5 sequence in one contiguous array, or may include the sequence as fragmented exons such as is often found for a flea gene. As used herein, the term “coding region” refers to a continuous linear array of nucleotides that translates into a protein. A full-length coding region is that coding region that is translated into a full-length, i.e., a complete protein as would be initially translated in its natural milieu, prior to any post-translational

10 modifications.

Table I represents a variety of flea HNC nucleic acid molecules of the present invention.

SEQ ID NO.	Name
1	2227-82HNC
2	2228-51HNC
3	2229-06HNC
4	2229-42HNC
5	2245-07HNC
6	2245-93HNC
7	2247-04HNC
8	2229-44HNC
9	2249-09HNC
10	2249-10HNC
11	2251-09HNC
12	2215-13HNC
13	2218-39HNC
14	2218-82HNC

Table II represents a variety of flea HMT nucleic acid molecules of the present invention.

SEQ ID NO:	Name		
15	2084-03HMT	56	2106-10HMT
16	2084-27HMT	57	2107-21HMT
17	2084-41HMT	58	2107-36HMT
18	2084-45HMT	59	2108-08HMT
19	2084-47HMT	60	2108-12HMT
20	2084-49HMT	61	2108-22HMT
21	2084-51HMT	62	2124-03HMT
22	2085-10HMT	63	2154-17HMT
23	2085-23HMT	64	2154-18HMT
24	2085-25HMT	65	2154-49HMT
25	2085-41HMT	66	2154-52HMT
26	2085-57HMT	67	2154-55HMT
27	2086-22HMT	68	2154-57HMT
28	2087-18HMT	69	2156-05HMT
29	2088-24HMT	70	2156-57HMT
30	2088-27HMT	71	2157-40HMT
31	2088-30HMT	72	2158-08HMT
32	2088-51HMT	73	2158-20HMT
33	2088-53HMT	74	2158-78HMT
34	2089-52HMT	75	2159-78HMT
35	2091-32HMT	76	2160-38HMT
36	2091-62HMT	77	2160-87HMT
37	2092-34HMT	78	2162-16HMT
38	2093-57HMT	79	2162-20HMT
39	2094-14HMT	80	2162-74HMT
40	2094-18HMT	81	2166-25HMT
41	2094-46HMT	82	2167-46HMT
42	2094-61HMT	83	2169-34HMT
43	2095-14HMT	84	2171-31HMT
44	2095-25HMT	85	2171-81HMT
45	2095-50HMT	86	2173-28HMT
46	2102-06HMT	87	2173-61HMT
47	2102-25HMT	88	2175-12HMT
48	2102-26HMT	89	2175-26HMT
49	2103-15HMT	90	2175-77HMT
50	2104-14HMT	91	2177-26HMT
51	2104-16HMT	92	2177-42HMT
52	2105-04HMT	93	2177-50HMT
53	2105-07HMT	94	2177-82HMT
54	2105-25HMT	95	2181-65HMT
55	2105-41HMT	96	2181-85HMT
		97	2183-34HMT

	SEQ ID NO	Name
	98	2183-77HMT
	99	2183-90HMT
	100	2187-45HMT
5	101	2188-28HMT
	102	2188-75HMT
	103	2191-26HMT
	104	2191-66HMT
	105	2192-23HMT
10	106	2192-42HMT
	107	2193-50HMT
	108	2194-30HMT
	109	2194-37HMT
	110	2195-34HMT
15	111	2195-80HMT
	112	2196-04HMT
	113	2196-29HMT
	114	2197-24HMT
	115	2202-47HMT
	116	2205-78HMT
20	117	2253-02HMT
	118	2253-05HMT
	119	2253-07HMT
	120	2253-21HMT
25	121	2253-73HMT
	122	2253-82HMT
	123	2253-83HMT
	124	2253-84HMT
	125	2253-96HMT
	126	2254-23HMT
30	127	2254-24HMT
	128	2254-29HMT
	129	2254-40HMT
	130	2254-53HMT
35	131	2254-57HMT
	132	2254-71HMT
	133	2254-85HMT
	134	2254-87HMT
	135	2254-92HMT
40	136	2255-14HMT
	137	2255-15HMT
	138	2255-20HMT
	139	2255-21HMT

140	2255-34HMT
141	2255-52HMT
142	2255-61HMT
143	2255-72HMT
144	2255-80HMT
145	2256-05HMT
146	2085-58HMT
147	2086-20HMT
148	2086-53HMT
149	2087-06HMT
150	2088-05HMT
151	2088-17HMT
152	2089-10HMT
153	2092-40HMT
154	2094-33HMT
155	2094-47HMT
156	2103-28HMT
157	2103-58HMT
158	2104-23HMT
159	2104-59HMT
160	2105-09HMT
161	2106-14HMT
162	2156-03HMT
163	2156-14HMT
164	2161-46HMT
165	2162-93HMT
166	2175-18HMT
167	2185-19HMT
168	2191-92HMT
169	2195-06HMT
170	2196-61HMT
171	2253-13HMT
172	2253-16HMT
173	2253-19HMT
174	2253-27HMT
175	2253-40HMT
176	2253-56HMT
177	2253-68HMT
178	2253-78HMT
179	2253-87HMT
180	2253-92HMT
181	2254-43HMT

	SEQ ID NO	Name
	182	2254-45HMT
	183	2254-51HMT
	184	2254-54HMT
5	185	2254-66HMT
	186	2254-88HMT
	187	2254-96HMT
	188	2255-17HMT
	189	2255-19HMT
10	190	2255-23HMT
	191	2255-25HMT
	192	2255-32HMT
	193	2255-39HMT
	194	2255-41HMT
	195	2255-45HMT
15	196	2255-60HMT
	197	2255-62HMT
	198	2255-67HMT
	199	2255-77HMT
	200	2255-79HMT
20	201	2255-82HMT
	202	2256-12HMT
	203	2256-22HMT
	204	2256-46HMT
	205	2256-54HMT
25	206	2256-58HMT
	207	2256-64HMT
	208	2229-13HMT
	209	2229-17HMT
	210	2229-20HMT
30	211	2229-31HMT
	212	2229-33HMT
	213	2229-34HMT
	214	2232-63HMT
	215	2232-70HMT
35	216	2232-81HMT
	217	2232-96HMT
	218	2233-03HMT
	219	2233-04HMT
	220	2233-09HMT
40	221	2233-14HMT
	222	2233-20HMT
	223	2233-33HMT

224	2233-38HMT
225	2233-39HMT
226	2233-48HMT
227	2233-54HMT
228	2233-56HMT
229	2233-60HMT
230	2233-64HMT
231	2233-70HMT
232	2233-71HMT
233	2233-75HMT
234	2233-76HMT
235	2233-77HMT
236	2233-92HMT
237	2233-95HMT
238	2234-08HMT
239	2234-15HMT
240	2234-17HMT
241	2234-18HMT
242	2234-30HMT
243	2234-33HMT
244	2234-35HMT
245	2234-37HMT
246	2234-57HMT
247	2234-65HMT
248	2240-04HMT
249	2241-07HMT
250	2241-08HMT
251	2241-09HMT
252	2229-71HMT
253	2229-86HMT
254	2230-04HMT
255	2230-09HMT
256	2230-12HMT
257	2230-13HMT
258	2230-30HMT
259	2230-31HMT
260	2230-36HMT
261	2230-40HMT
262	2230-42HMT
263	2230-44HMT
264	2230-45HMT
265	2230-61HMT

	SEQ ID NO.	Name
	266	2230-63HMT
	267	2230-68HMT
	268	2230-70HMT
	269	2230-72HMT
5	270	2230-76HMT
	271	2230-78HMT
	272	2230-86HMT
	273	2230-92HMT
10	274	2231-05HMT
	275	2231-06HMT
	276	2231-09HMT
	277	2231-12HMT
	278	2231-17HMT
	279	2231-30HMT
15	280	2231-34HMT
	281	2231-45HMT
	282	2231-96HMT
	283	2232-07HMT
	284	2232-09HMT
20	285	2232-13HMT
	286	2232-23HMT
	287	2232-31HMT
	288	2232-35HMT
	289	2232-40HMT
25	290	2232-46HMT
	291	2232-54HMT
	292	2229-51HMT
	293	2230-60HMT
	294	2230-81HMT
30	295	2231-03HMT
	296	2231-24HMT
	297	2231-56HMT
	298	2233-59HMT
	299	2234-12HMT
35	300	2234-76HMT
	301	2234-86HMT
	302	2240-11HMT
	303	2240-14HMT
	304	2240-17HMT
40	305	2240-23HMT
	306	2240-26HMT
	307	2240-28HMT

308	2240-31HMT
309	2240-44HMT
310	2240-63HMT
311	2240-66HMT
312	2240-70HMT
313	2240-72HMT
314	2240-94HMT
315	2241-10HMT
316	2241-12HMT
317	2241-40HMT
318	2241-54HMT
319	2241-56HMT
320	2241-60HMT
321	2241-66HMT
322	2241-74HMT
323	2241-82HMT
324	2241-83HMT
325	2241-86HMT
326	2241-87HMT
327	2243-10HMT
328	2243-18HMT
329	2243-20HMT
330	2243-22HMT
331	2243-27HMT
332	2243-28HMT
333	2243-31HMT
334	2243-32HMT
335	2243-38HMT
336	2243-41HMT
337	2243-45HMT
338	2243-48HMT
339	2243-52HMT
340	2243-73HMT
341	2243-86HMT
342	2243-87HMT
343	2243-94HMT
344	2244-12HMT
345	2244-25HMT
346	2244-29HMT
347	2244-44HMT
348	2244-54HMT
349	2244-71HMT

	SEQ ID NO:	Name
	350	2244-75HMT
	351	2084-19
	352	2084-25
5	353	2084-57
	354	2084-58
	355	2084-64
	356	2085-06
	357	2085-12
10	358	2085-27
	359	2085-28
	360	2086-08
	361	2086-14
	362	2086-16
	363	2086-21
15	364	2086-23
	365	2086-46
	366	2086-51
	367	2086-56
	368	2086-57
20	369	2086-59
	370	2086-62
	371	2087-05
	372	2087-16
	373	2087-19
25	374	2087-24
	375	2087-53
	376	2087-55
	377	2088-06
	378	2088-08
30	379	2088-12
	380	2088-13
	381	2088-14
	382	2088-16
	383	2088-18
35	384	2088-23
	385	2088-28
	386	2088-34
	387	2088-42
	388	2088-44
40	389	2088-45
	390	2088-50
	391	2089-04
	392	2089-15
	393	2089-16

394	2089-17
395	2089-18
396	2089-19
397	2089-24
398	2089-31
399	2089-38
400	2089-40
401	2089-41
402	2089-53
403	2089-58
404	2090-02
405	2090-12
406	2090-16
407	2090-29
408	2090-37
409	2090-39
410	2090-41
411	2090-50
412	2090-52
413	2090-56
414	2091-09
415	2091-15
416	2091-16
417	2091-18
418	2091-24
419	2091-25
420	2091-28
421	2091-52
422	2092-02
423	209206-
424	2092-08
425	2092-15
426	2092-22
427	2092-24
428	2092-26
429	2092-30
430	2092-35
431	2092-53
432	2092-55
433	2092-63
434	2093-11
435	2093-19
436	2093-24
437	2093-27

	SEQ ID NO:	Name
	438	2093-28
	439	2093-33
	440	2093-36
5	441	2093-37
	442	2093-38
	443	2093-40
	444	2093-51
	445	2093-52
10	446	2093-53
	447	2093-55
	448	2093-56
	449	2093-60
	450	2094-02
	451	2094-04
15	452	2094-06
	453	2094-09
	454	2094-10
	455	2094-13
	456	2094-15
20	457	2094-23
	458	2094-27
	459	2094-29
	460	2094-34
	461	2094-37
25	462	2094-39
	463	2094-41
	464	2094-45
	465	2094-49
	466	2094-51
30	467	2094-53
	468	2094-54
	469	2094-59
	470	2095-02
	471	2095-07
35	472	2095-16
	473	2095-19
	474	2095-21
	475	2095-24
	476	2095-32
40	477	2095-35
	478	2095-37
	479	2095-39
	480	2095-42
	481	2095-44

	482	2095-46
	483	2095-49
	484	2095-53
	485	2095-56
	486	2095-58
	487	2095-61
	488	2095-62
	489	2095-64
	490	2102-03
	491	2102-04
	492	2102-05
	493	2102-07
	494	2102-08
	495	2102-10
	496	2102-14
	497	2102-15
	498	2102-31
	499	2102-34
	500	2102-39
	501	2102-40
	502	2102-42
	503	2102-43
	504	2102-52
	505	2102-56
	506	2102-57
	507	2102-62
	508	2103-04
	509	2103-06
	510	2103-07
	511	2103-10
	512	2103-16
	513	2103-18
	514	2103-19
	515	2103-20
	516	2103-21
	517	2103-22
	518	2103-23
	519	2103-25
	520	2103-26
	521	2103-32
	522	2103-34
	523	2103-36
	524	2103-38
	525	2103-39

	SEQ ID NO:	Name
	526	2103-42
	527	2103-43
	528	2103-46
	529	2103-51
5	530	2103-52
	531	2103-54
	532	2103-59
	533	2103-64
	534	2104-02
10	535	2104-09
	536	2104-11
	537	2104-12
	538	2104-17
	539	2104-19
15	540	2104-27
	541	2104-30
	542	2104-33
	543	2104-36
	544	2104-37
20	545	2104-39
	546	2104-42
	547	2104-49
	548	2104-52
	549	2104-53
25	550	2104-57
	551	2104-62
	552	2105-03
	553	2105-05
	554	2105-06
30	555	2105-08
	556	2105-13
	557	2105-22
	558	2105-24
	559	2105-26
35	560	2105-27
	561	2105-29
	562	2105-30
	563	2105-36
	564	2105-39
40	565	2105-47
	566	2105-49
	567	2105-52
	568	2105-58
	569	2105-59

570	2105-60
571	2106-03
572	2106-05
573	2106-06
574	2106-08
575	2106-13
576	2106-22
577	2106-24
578	2106-26
579	2106-27
580	2106-29
581	2106-30
582	2106-36
583	2106-39
584	2106-47
585	2106-49
586	2106-52
587	2106-58
588	2106-59
589	2106-60
590	2107-05
591	2107-07
592	2107-08
593	2107-14
594	2107-20
595	2107-23
596	2107-27
597	2107-31
598	2107-33
599	2107-34
600	2107-35
601	2107-42
602	2107-44
603	2107-47
604	2107-48
605	2107-49
606	2107-51
607	2107-53
608	2107-56
609	2107-60
610	2107-62
611	2107-63
612	2108-15
613	2108-16



	SEQ ID NO:	Name
	614	2108-18
	615	2108-20
	616	2108-24
5	617	2108-25
	618	2108-30
	619	2108-32
	620	2108-33
	621	2108-34
10	622	2108-43
	623	2108-45
	624	2108-50
	625	2108-55
	626	2108-61
	627	2108-62
15	628	2109-03
	629	2109-05
	630	2109-09
	631	2109-11
	632	2109-12
20	633	2109-14
	634	2109-15
	635	2109-17
	636	2109-20
	637	2109-22
25	638	2109-23
	639	2109-25
	640	2109-27
	641	2109-29
	642	2109-32
30	643	2109-33
	644	2109-34
	645	2109-36
	646	2109-41
	647	2109-43
35	648	2109-45
	649	2109-46
	650	2109-48
	651	2109-49
	652	2109-50
40	653	2109-57
	654	2109-60
	655	2109-61
	656	2109-62
	657	2109-63

658	2154-02
659	2154-03
660	2154-05
661	2154-07
662	2154-12
663	2154-13
664	2154-14
665	2154-15
666	2154-19
667	2154-22
668	2154-26
669	2154-35
670	2154-37
671	2154-39
672	2154-40
673	2154-44
674	2154-53
675	2154-59
676	2154-62
677	2154-63
678	2154-67
679	2154-69
680	2154-88
681	2156-04
682	2156-08
683	2156-13
684	2156-16
685	2156-19
686	2156-20
687	2156-22
688	2156-24
689	2156-25
690	2156-30
691	2156-31
692	2156-32
693	2156-33
694	2156-35
695	2156-39
696	2156-45
697	2156-46
698	2156-49
699	2156-55
700	2156-61
701	2156-64

	SEQ ID NO:	Name
	702	2157-03
	703	2157-04
	704	2157-07
5	705	2157-09
	706	2157-13
	707	2157-17
	708	2157-18
	709	2157-19
10	710	2157-22
	711	2157-23
	712	2157-24
	713	2157-25
	714	2157-26
15	715	2157-28
	716	2157-35
	717	2157-37
	718	2157-38
	719	2157-41
20	720	2157-42
	721	2157-46
	722	2157-49
	723	2157-50
	724	2157-52
25	725	2157-57
	726	2157-63
	727	2157-65
	728	2157-71
	729	2157-72
30	730	2157-74
	731	2157-75
	732	2157-80
	733	2157-93
	734	2158-08
35	735	2158-10
	736	2158-13
	737	2158-15
	738	2158-16
	739	2158-20
40	740	2158-26
	741	2158-32
	742	2158-38
	743	2158-40
	744	2158-43
	745	2158-44

746	2158-45
747	2158-46
748	2159-16
749	2159-18
750	2159-24
751	2159-26
752	2159-28
753	2159-33
754	2159-37
755	2159-38
756	2159-40
757	2159-42
758	2159-43
759	2159-44
760	2159-45
761	2159-46
762	2159-50
763	2159-52
764	2159-61
765	2159-62
766	2159-65
767	2159-67
768	2159-70
769	2159-71
770	2159-72
771	2159-73
772	2159-74
773	2159-80
774	2159-83
775	2159-86
776	2159-89
777	2159-90
778	2159-93
779	2159-95
780	2160-04
781	2160-05
782	2160-08
783	2160-10
784	2160-15
785	2160-20
786	2160-22
787	2160-24
788	2160-33
789	2160-37

	SEQ ID NO:	Name:
	790	2160-40
	791	2160-41
	792	2160-43
	793	2160-46
5	794	2160-51
	795	2160-52
	796	2160-56
	797	2160-57
	798	2160-64
10	799	2160-66
	800	2160-72
	801	2160-73
	802	2160-76
	803	2160-78
15	804	2160-92
	805	2160-94
	806	2160-95
	807	2160-96
	808	2161-02
20	809	2161-22
	810	2161-24
	811	2161-25
	812	2161-26
	813	2161-27
25	814	2161-29
	815	2161-30
	816	2161-31
	817	2161-32
	818	2161-33
30	819	2161-36
	820	2161-41
	821	2161-42
	822	2161-50
	823	2161-51
35	824	2161-53
	825	2161-55
	826	2161-61
	827	2161-63
	828	2161-66
40	829	2161-68
	830	2161-71
	831	2161-73
	832	2161-84
	833	2162-04

834	2162-07
835	2162-09
836	2162-10
837	2162-14
838	2162-15
839	2162-17
840	2162-19
841	2162-22
842	2162-23
843	2162-27
844	2162-29
845	2162-37
846	2162-39
847	2162-43
848	2162-51
849	2162-52
850	2162-54
851	2162-55
852	2162-56
853	2162-57
854	2162-58
855	2162-60
856	2162-61
857	2162-65
858	2162-67
859	2162-70
860	2162-76
861	2162-77
862	2162-79
863	2162-82
864	2162-84
865	2162-85
866	2162-86
867	2162-89
868	2162-90
869	2162-94
870	2163-05
871	2163-06
872	2163-08
873	2163-09
874	2163-10
875	2163-12
876	2163-13
877	2163-14

	SEQ ID NO.	Name
	878	2163-15
	879	2163-16
	880	2163-17
5	881	2163-19
	882	2163-20
	883	2163-30
	884	2163-33
	885	2163-34
10	886	2163-35
	887	2163-37
	888	2163-38
	889	2163-39
	890	2163-40
	891	2163-41
15	892	2163-46
	893	2163-47
	894	2163-59
	895	2163-63
	896	2163-66
20	897	2163-67
	898	2163-68
	899	2163-69
	900	2163-70
	901	2163-72
25	902	2163-74
	903	2163-75
	904	2163-80
	905	2163-88
	906	2163-96
30	907	2164-02
	908	2164-03
	909	2164-04
	910	2164-05
	911	2164-06
35	912	2164-09
	913	2164-10
	914	2164-11
	915	2164-12
40	916	2164-14
	917	2164-15
	918	2164-17
	919	2164-20
	920	2164-21
	921	2164-22

922	2164-23
923	2164-26
924	2164-27
925	2164-28
926	2164-29
927	2164-30
928	2164-31
929	2164-32
930	2164-33
931	2164-34
932	2164-36
933	2164-37
934	2164-41
935	2164-42
936	2164-44
937	2164-46
938	2164-48
939	2164-49
940	2164-50
941	2164-51
942	2164-52
943	2164-53
944	2164-54
945	2164-56
946	2164-57
947	2164-58
948	2164-59
949	2164-61
950	2164-62
951	2164-68
952	2164-69
953	2164-70
954	2164-71
955	2164-72
956	2164-73
957	2164-74
958	2164-75
959	2164-77
960	2164-78
961	2164-79
962	2164-80
963	2164-81
964	2164-83
965	2164-84

	SEQ ID NO:	Name
	966	2164-85
	967	2164-86
	968	2164-87
	969	2164-88
5	970	2164-89
	971	2164-91
	972	2164-92
	973	2164-93
	974	2164-94
10	975	2164-95
	976	2164-96
	977	2165-02
	978	2165-03
	979	2165-07
15	980	2165-09
	981	2165-10
	982	2165-11
	983	2165-13
	984	2165-15
20	985	2165-19
	986	2165-22
	987	2165-23
	988	2165-25
	989	2165-29
25	990	2165-34
	991	2165-38
	992	2165-39
	993	2165-40
	994	2165-43
30	995	2165-44
	996	2165-46
	997	2165-49
	998	2165-50
	999	2165-51
35	1000	2165-52
	1001	2165-53
	1002	2165-55
	1003	2165-61
	1004	2165-62
40	1005	2165-71
	1006	2165-72
	1007	2165-73
	1008	2165-75
	1009	2165-77

1010	2165-78
1011	2165-79
1012	2165-80
1013	2165-81
1014	2165-91
1015	2165-92
1016	2165-93
1017	2165-95
1018	2165-96
1019	2166-04
1020	2166-06
1021	2166-07
1022	2166-08
1023	2166-13
1024	2166-14
1025	2166-16
1026	2166-18
1027	2166-20
1028	2166-30
1029	2166-34
1030	2166-36
1031	2166-37
1032	2166-38
1033	2166-39
1034	2166-43
1035	2166-45
1036	2166-48
1037	2166-49
1038	2166-52
1039	2166-54
1040	2166-55
1041	2166-57
1042	2166-59
1043	2166-63
1044	2166-66
1045	2166-67
1046	2166-68
1047	2166-69
1048	2166-70
1049	2166-72
1050	2166-74
1051	2166-85
1052	2166-86
1053	2167-02

	SEQ ID NO:	Name
	1054	2167-06
	1055	2167-08
	1056	2167-11
5	1057	2167-12
	1058	2167-17
	1059	2167-19
	1060	2167-21
	1061	2167-22
	1062	2167-23
10	1063	2167-24
	1064	2167-27
	1065	2167-29
	1066	2167-30
	1067	2167-32
15	1068	2167-34
	1069	2167-37
	1070	2167-38
	1071	2167-39
	1072	2167-41
20	1073	2167-43
	1074	2167-50
	1075	2167-54
	1076	2167-55
	1077	2167-57
25	1078	2167-58
	1079	2167-64
	1080	2167-65
	1081	2167-71
	1082	2167-72
30	1083	2167-73
	1084	2167-74
	1085	2167-75
	1086	2167-76
	1087	2167-77
35	1088	2167-81
	1089	2167-83
	1090	2167-86
	1091	2167-89
	1092	2167-93
40	1093	2167-94
	1094	2167-95
	1095	2169-02
	1096	2169-03
	1097	2169-11

1098	2169-15
1099	2169-18
1100	2169-20
1101	2169-21
1102	2169-23
1103	2169-25
1104	2169-28
1105	2169-29
1106	2169-31
1107	2169-36
1108	2169-37
1109	2169-38
1110	2169-45
1111	2169-48
1112	2169-50
1113	2169-53
1114	2169-54
1115	2169-55
1116	2169-56
1117	2169-58
1118	2169-59
1119	2169-60
1120	2169-61
1121	2169-62
1122	2169-63
1123	2169-64
1124	2169-67
1125	2169-68
1126	2169-72
1127	2169-73
1128	2169-75
1129	2169-76
1130	2169-77
1131	2169-83
1132	2169-84
1133	2169-87
1134	2169-89
1135	2169-90
1136	2169-92
1137	2169-93
1138	2169-94
1139	2171-02
1140	2171-05
1141	2171-07

	SEQ ID NO.	Name
	1142	2171-08
	1143	2171-10
	1144	2171-12
5	1145	2171-14
	1146	2171-15
	1147	2171-18
	1148	2171-19
	1149	2171-25
	1150	2171-26
10	1151	2171-27
	1152	2171-28
	1153	2171-29
	1154	2171-36
	1155	2171-38
15	1156	2171-42
	1157	2171-44
	1158	2171-45
	1159	2171-48
	1160	2171-49
20	1161	2171-53
	1162	2171-58
	1163	2171-59
	1164	2171-62
	1165	2171-65
25	1166	2171-67
	1167	2171-70
	1168	2171-71
	1169	2171-74
	1170	2171-75
30	1171	2171-77
	1172	2171-78
	1173	2171-83
	1174	2171-86
	1175	2171-87
35	1176	2171-90
	1177	2171-91
	1178	2171-94
	1179	2171-95
	1180	2171-96
40	1181	2173-03
	1182	2173-04
	1183	2173-05
	1184	2173-06
	1185	2173-15

1186	2173-17
1187	2173-20
1188	2173-24
1189	2173-26
1190	2173-27
1191	2173-30
1192	2173-31
1193	2173-33
1194	2173-35
1195	2173-37
1196	2173-41
1197	2173-46
1198	2173-49
1199	2173-60
1200	2173-62
1201	2173-65
1202	2173-66
1203	2173-69
1204	2173-72
1205	2173-73
1206	2173-77
1207	2173-79
1208	2173-80
1209	2173-82
1210	2173-87
1211	2173-88
1212	2173-89
1213	2173-92
1214	2175-03
1215	2175-05
1216	2175-09
1217	2175-10
1218	2175-11
1219	2175-14
1220	2175-16
1221	2175-17
1222	2175-21
1223	2175-22
1224	2175-23
1225	2175-27
1226	2175-29
1227	2175-30
1228	2175-31
1229	2175-33

	SEQ ID NO:	Name
	1230	2175-35
	1231	2175-36
	1232	2175-40
5	1233	2175-41
	1234	2175-42
	1235	2175-43
	1236	2175-47
	1237	2175-50
	1238	2175-51
10	1239	2175-52
	1240	2175-53
	1241	2175-54
	1242	2175-55
	1243	2175-56
15	1244	2175-57
	1245	2175-59
	1246	2175-60
	1247	2175-61
	1248	2175-63
20	1249	2175-65
	1250	2175-66
	1251	2175-67
	1252	2175-68
	1253	2175-69
25	1254	2175-70
	1255	2175-79
	1256	2175-80
	1257	2175-81
	1258	2175-85
30	1259	2175-87
	1260	2175-93
	1261	2175-94
	1262	2175-95
	1263	2177-02
35	1264	2177-03
	1265	2177-04
	1266	2177-05
	1267	2177-09
	1268	2177-25
40	1269	2177-28
	1270	2177-30
	1271	2177-32
	1272	2177-35
	1273	2177-36

1274	2177-37
1275	2177-38
1276	2177-43
1277	2177-49
1278	2177-55
1279	2177-56
1280	2177-58
1281	2177-62
1282	2177-63
1283	2177-65
1284	2177-66
1285	2177-67
1286	2177-68
1287	2177-71
1288	2177-73
1289	2177-75
1290	2177-76
1291	2177-77
1292	2177-83
1293	2177-84
1294	2177-85
1295	2177-87
1296	2177-88
1297	2177-89
1298	2177-90
1299	2177-93
1300	2177-95
1301	2179-04
1302	2179-06
1303	2179-08
1304	2179-13
1305	2179-14
1306	2179-15
1307	2179-16
1308	2179-17
1309	2179-21
1310	2179-25
1311	2179-26
1312	2179-31
1313	2179-33
1314	2179-34
1315	2179-36
1316	2179-37
1317	2179-38



	SEQ ID NO:	Name
	1318	2179-40
	1319	2179-41
	1320	2179-45
5	1321	2179-50
	1322	2179-55
	1323	2179-57
	1324	2179-58
	1325	2179-60
10	1326	2179-61
	1327	2179-64
	1328	2179-65
	1329	2179-66
	1330	2179-67
	1331	2179-68
15	1332	2179-70
	1333	2179-71
	1334	2179-72
	1335	2179-74
	1336	2179-75
20	1337	2179-76
	1338	2179-78
	1339	2179-80
	1340	2179-81
	1341	2179-82
25	1342	2179-84
	1343	2179-85
	1344	2179-89
	1345	2179-90
	1346	2179-91
30	1347	2179-92
	1348	2179-93
	1349	2179-95
	1350	2179-96
	1351	2181-03
35	1352	2181-06
	1353	2181-07
	1354	2181-08
	1355	2181-11
	1356	2181-13
40	1357	2181-15
	1358	2181-16
	1359	2181-17
	1360	2181-20
	1361	2181-21

1362	2181-25
1363	2181-28
1364	2181-30
1365	2181-31
1366	2181-32
1367	2181-33
1368	2181-34
1369	2181-38
1370	2181-40
1371	2181-41
1372	2181-43
1373	2181-44
1374	2181-47
1375	2181-48
1376	2181-49
1377	2181-53
1378	2181-54
1379	2181-55
1380	2181-57
1381	2181-58
1382	2181-59
1383	2181-60
1384	2181-61
1385	2181-62
1386	2181-63
1387	2181-67
1388	2181-70
1389	2181-71
1390	2181-72
1391	2181-74
1392	2181-77
1393	2181-78
1394	2181-79
1395	2181-80
1396	2181-83
1397	2181-85
1398	2181-86
1399	2181-88
1400	2181-89
1401	2181-90
1402	2181-95
1403	2181-96
1404	2183-02
1405	2183-03

	SEQ ID NO:	Name
	1406	2183-04
	1407	2183-09
	1408	2183-12
	1409	2183-18
5	1410	2183-19
	1411	2183-23
	1412	2183-26
	1413	2183-27
	1414	2183-29
10	1415	2183-31
	1416	2183-34
	1417	2183-35
	1418	2183-37
	1419	2183-42
15	1420	2183-43
	1421	2183-47
	1422	2183-48
	1423	2183-54
	1424	2183-56
20	1425	2183-57
	1426	2183-60
	1427	2183-63
	1428	2183-64
	1429	2183-66
25	1430	2183-67
	1431	2183-72
	1432	2183-73
	1433	2183-74
	1434	2183-75
30	1435	2183-76
	1436	2183-78
	1437	2183-79
	1438	2183-80
	1439	2183-82
35	1440	2183-83
	1441	2183-84
	1442	2183-87
	1443	2183-89
	1444	2183-91
40	1445	2183-92
	1446	2183-94
	1447	2185-02
	1448	2185-03
	1449	2185-06

1450	2185-07
1451	2185-08
1452	2185-11
1453	2185-13
1454	2185-14
1455	2185-17
1456	2185-18
1457	2185-25
1458	2185-26
1459	2185-30
1460	2185-31
1461	2185-32
1462	2185-35
1463	2185-37
1464	2185-39
1465	2185-40
1466	2185-42
1467	2185-45
1468	2185-46
1469	2185-47
1470	2185-50
1471	2185-58
1472	2185-63
1473	2185-64
1474	2185-65
1475	2185-67
1476	2185-68
1477	2185-72
1478	2185-73
1479	2185-74
1480	2185-76
1481	2185-77
1482	2185-79
1483	2185-83
1484	2185-85
1485	2185-86
1486	2185-87
1487	2185-89
1488	2185-90
1489	2185-92
1490	2185-93
1491	2185-94
1492	2185-96
1493	2187-02

	SEQ ID NO:	Name
	1494	2187-06
	1495	2187-08
	1496	2187-11
5	1497	2187-12
	1498	2187-15
	1499	2187-16
	1500	2187-19
	1501	2187-22
10	1502	2187-26
	1503	2187-27
	1504	2187-28
	1505	2187-29
	1506	2187-30
	1507	2187-34
15	1508	2187-35
	1509	2187-37
	1510	2187-38
	1511	2187-39
	1512	2187-40
20	1513	2187-41
	1514	2187-42
	1515	2187-44
	1516	2187-46
	1517	2187-48
25	1518	2187-50
	1519	2187-51
	1520	2187-52
	1521	2187-54
	1522	2187-55
30	1523	2187-56
	1524	2187-57
	1525	2187-58
	1526	2187-60
	1527	2187-61
35	1528	2187-62
	1529	2187-63
	1530	2187-70
	1531	2187-71
	1532	2187-72
40	1533	2187-73
	1534	2187-80
	1535	2187-84
	1536	2187-85
	1537	2187-86

1538	2187-87
1539	2187-88
1540	2187-91
1541	2187-94
1542	2187-96
1543	2188-04
1544	2188-08
1545	2188-09
1546	2188-11
1547	2188-18
1548	2188-20
1549	2188-21
1550	2188-23
1551	2188-25
1552	2188-31
1553	2188-37
1554	2188-38
1555	2188-42
1556	2188-43
1557	2188-44
1558	2188-47
1559	2188-50
1560	2188-53
1561	2188-55
1562	2188-57
1563	2188-59
1564	2188-61
1565	2188-64
1566	2188-69
1567	2188-70
1568	2188-71
1569	2188-73
1570	2188-76
1571	2188-79
1572	2188-84
1573	2188-85
1574	2188-87
1575	2188-88
1576	2188-89
1577	2188-94
1578	2188-95
1579	2188-96
1580	2189-02
1581	2189-03

	SEQ ID NO:	Name:
	1582	2189-04
	1583	2189-05
	1584	2189-06
5	1585	2189-07
	1586	2189-09
	1587	2189-10
	1588	2189-11
	1589	2189-12
	1590	2189-14
10	1591	2189-15
	1592	2189-16
	1593	2189-18
	1594	2189-19
	1595	2189-20
15	1596	2189-22
	1597	2189-23
	1598	2189-24
	1599	2189-25
	1600	2189-26
20	1601	2189-27
	1602	2189-28
	1603	2189-29
	1604	2189-30
	1605	2189-32
25	1606	2189-33
	1607	2189-34
	1608	2189-35
	1609	2189-36
	1610	2189-37
30	1611	2189-38
	1612	2189-40
	1613	2189-41
	1614	2189-43
	1615	2189-44
35	1616	2189-46
	1617	2189-47
	1618	2189-49
	1619	2189-50
	1620	2189-52
40	1621	2189-54
	1622	2189-58
	1623	2189-59
	1624	2189-60
	1625	2189-61

1626	2189-62
1627	2189-64
1628	2189-68
1629	2189-72
1630	2189-77
1631	2189-78
1632	2189-79
1633	2189-80
1634	2189-82
1635	2189-88
1636	2189-93
1637	2189-95
1638	2191-05
1639	2191-06
1640	2191-10
1641	2191-11
1642	2191-13
1643	2191-14
1644	2191-15
1645	2191-16
1646	2191-19
1647	2191-21
1648	2191-26
1649	2191-30
1650	2191-31
1651	2191-32
1652	2191-36
1653	2191-37
1654	2191-39
1655	2191-40
1656	2191-42
1657	2191-43
1658	2191-45
1659	2191-48
1660	2191-49
1661	2191-50
1662	2191-54
1663	2191-55
1664	2191-56
1665	2191-61
1666	2191-62
1667	2191-64
1668	2191-65
1669	2191-70

	SEQ ID NO:	Name
	1670	2191-71
	1671	2191-74
	1672	2191-75
5	1673	2191-80
	1674	2191-84
	1675	2191-86
	1676	2191-89
	1677	2191-91
	1678	2191-93
10	1679	2191-95
	1680	2192-04
	1681	2192-05
	1682	2192-10
	1683	2192-11
15	1684	2192-13
	1685	2192-16
	1686	2192-18
	1687	2192-19
	1688	2192-20
20	1689	2192-21
	1690	2192-24
	1691	2192-27
	1692	2192-32
	1693	2192-33
25	1694	2192-34
	1695	2192-39
	1696	2192-40
	1697	2192-41
	1698	2192-45
30	1699	2192-48
	1700	2192-49
	1701	2192-51
	1702	2192-52
	1703	2192-54
35	1704	2192-55
	1705	2192-56
	1706	2192-57
	1707	2192-58
	1708	2192-61
40	1709	2192-62
	1710	2192-64
	1711	2192-65
	1712	2192-67
	1713	2192-73

1714	2192-75
1715	2192-76
1716	2192-80
1717	2192-81
1718	2192-83
1719	2192-84
1720	2192-87
1721	2192-89
1722	2192-90
1723	2192-92
1724	2192-93
1725	2192-95
1726	2193-02
1727	2193-05
1728	2193-06
1729	2193-07
1730	2193-08
1731	2193-09
1732	2193-11
1733	2193-12
1734	2193-13
1735	2193-14
1736	2193-15
1737	2193-16
1738	2193-19
1739	2193-20
1740	2193-21
1741	2193-22
1742	2193-23
1743	2193-26
1744	2193-28
1745	2193-30
1746	2193-32
1747	2193-33
1748	2193-34
1749	2193-35
1750	2193-38
1751	2193-40
1752	2193-41
1753	2193-42
1754	2193-44
1755	2193-46
1756	2193-48
1757	2193-55

	SEQ ID NO.	Name
	1758	2193-57
	1759	2193-58
	1760	2193-59
5	1761	2193-62
	1762	2193-63
	1763	2193-64
	1764	2193-66
	1765	2193-68
	1766	2193-69
10	1767	2193-70
	1768	2193-72
	1769	2193-73
	1770	2193-74
	1771	2193-76
15	1772	2193-78
	1773	2193-81
	1774	2193-82
	1775	2193-83
	1776	2193-84
20	1777	2193-85
	1778	2193-86
	1779	2193-87
	1780	2193-88
	1781	2193-91
25	1782	2194-02
	1783	2194-03
	1784	2194-04
	1785	2194-05
	1786	2194-06
30	1787	2194-08
	1788	2194-10
	1789	2194-11
	1790	2194-12
	1791	2194-15
35	1792	2194-19
	1793	2194-21
	1794	2194-25
	1795	2194-26
	1796	2194-31
40	1797	2194-33
	1798	2194-34
	1799	2194-36
	1800	2194-44
	1801	2194-45

1802	2194-46
1803	2194-47
1804	2194-48
1805	2194-49
1806	2194-53
1807	2194-54
1808	2194-56
1809	2194-57
1810	2194-58
1811	2194-59
1812	2194-60
1813	2194-61
1814	2194-64
1815	2194-65
1816	2194-66
1817	2194-67
1818	2194-68
1819	2194-70
1820	2194-71
1821	2194-73
1822	2194-74
1823	2194-76
1824	2194-77
1825	2194-79
1826	2194-80
1827	2194-82
1828	2194-87
1829	2194-89
1830	2194-90
1831	2194-91
1832	2194-94
1833	2195-03
1834	2195-04
1835	2195-13
1836	2195-18
1837	2195-19
1838	2195-20
1839	2195-21
1840	2195-23
1841	2195-25
1842	2195-27
1843	2195-29
1844	2195-32
1845	2195-33

	SEQ ID NO:	Name
	1846	2195-35
	1847	2195-36
	1848	2195-37
	1849	2195-38
5	1850	2195-39
	1851	2195-41
	1852	2195-42
	1853	2195-43
	1854	2195-44
10	1855	2195-46
	1856	2195-48
	1857	2195-50
	1858	2195-52
	1859	2195-56
15	1860	2195-58
	1861	2195-59
	1862	2195-62
	1863	2195-63
	1864	2195-64
20	1865	2195-65
	1866	2195-68
	1867	2195-70
	1868	2195-71
	1869	2195-74
25	1870	2195-75
	1871	2195-77
	1872	2195-78
	1873	2195-79
	1874	2195-81
30	1875	2195-82
	1876	2195-87
	1877	2195-88
	1878	2195-89
	1879	2195-90
35	1880	2195-91
	1881	2195-93
	1882	2195-96
	1883	2196-02
	1884	2196-05
40	1885	2196-06
	1886	2196-08
	1887	2196-09
	1888	2196-11
	1889	2196-15

1890	2196-16
1891	2196-17
1892	2196-23
1893	2196-25
1894	2196-28
1895	2196-32
1896	2196-33
1897	2196-34
1898	2196-35
1899	2196-36
1900	2196-38
1901	2196-39
1902	2196-40
1903	2196-41
1904	2196-45
1905	2196-49
1906	2196-51
1907	2196-55
1908	2196-56
1909	2196-57
1910	2196-58
1911	2196-66
1912	2196-67
1913	2196-68
1914	2196-73
1915	2196-77
1916	2196-80
1917	2196-87
1918	2196-89
1919	2196-91
1920	2196-92
1921	2196-93
1922	2196-94
1923	2196-96
1924	2197-02
1925	2197-04
1926	2197-05
1927	2197-06
1928	2197-07
1929	2197-09
1930	2197-10
1931	2197-11
1932	2197-12
1933	2197-13

	SEQ ID NO:	Name
	1934	2197-15
	1935	2197-16
	1936	2197-19
5	1937	2197-20
	1938	2197-27
	1939	2197-29
	1940	2197-30
	1941	2197-31
10	1942	2197-35
	1943	2197-36
	1944	2197-37
	1945	2197-38
	1946	2197-39
15	1947	2197-40
	1948	2197-41
	1949	2197-43
	1950	2197-44
	1951	2197-45
20	1952	2197-48
	1953	2197-49
	1954	2197-52
	1955	2197-55
	1956	2197-56
25	1957	2197-57
	1958	2197-61
	1959	2197-62
	1960	2197-63
	1961	2197-65
30	1962	2197-66
	1963	2197-67
	1964	2197-69
	1965	2197-71
	1966	2197-74
35	1967	2197-75
	1968	2197-76
	1969	2197-77
	1970	2197-79
	1971	2197-82
40	1972	2197-83
	1973	2197-84
	1974	2197-85
	1975	2197-87
	1976	2197-88
	1977	2197-89

1978	2197-94
1979	2197-96
1980	2202-18
1981	2202-19
1982	2202-22
1983	2202-25
1984	2202-28
1985	2202-30
1986	2202-31
1987	2202-32
1988	2202-38
1989	2202-44
1990	2202-46
1991	2202-55
1992	2202-64
1993	2202-66
1994	2202-67
1995	2202-68
1996	2202-71
1997	2202-72
1998	2202-73
1999	2202-04
2000	2202-87
2001	2202-93
2002	2202-94
2003	2203-12
2004	2203-20
2005	2203-24
2006	2202-26
2007	2202-30
2008	2202-52
2009	2202-74
2010	2202-78
2011	2202-87
2012	2202-93
2013	2202-94
2014	2204-02
2015	2204-08
2016	2204-12
2017	2204-14
2018	2204-42
2019	2204-43
2020	2204-49
2021	2204-51



SEQ ID NO:	Name
2022	2204-53
2023	2204-54
2024	2204-59
2025	2204-60
2026	2204-61
2027	2204-70
2028	2204-71
2029	2204-73
2030	2204-74
2031	2204-78
2032	2204-83
2033	2204-90
2034	2205-05
2035	2205-11
2036	2205-13
2037	2205-21
2038	2205-22
2039	2205-39
2040	2205-42
2041	2205-44
2042	2205-81
2043	2205-88
2044	2205-94
2045	2229-15
2046	2229-21
2047	2229-23
2048	2229-26
2049	2229-28
2050	2229-30
2051	2229-32
2052	2229-43
2053	2229-52
2054	2229-66
2055	2229-85
2056	2230-17
2057	2230-18
2058	2230-25
2059	2230-29
2060	2230-32
2061	2230-34
2062	2230-46
2063	2230-47
2064	2230-58
2065	2230-65

2066	2230-73
2067	2230-74
2068	2230-95
2069	2231-04
2070	2231-13
2071	2231-16
2072	2231-21
2073	2231-25
2074	2231-28
2075	2231-48
2076	2231-54
2077	2231-58
2078	2231-61
2079	2231-69
2080	2231-93
2081	2232-17
2082	2232-20
2083	2232-26
2084	2232-29
2085	2232-36
2086	2232-37
2087	2232-45
2088	2232-47
2089	2232-48
2090	2232-53
2091	2232-62
2092	2232-75
2093	2232-76
2094	2232-88
2095	2232-89
2096	2232-90
2097	2232-92
2098	2232-93
2099	2232-94
2100	2233-07
2101	2233-15
2102	2233-17
2103	2233-23
2104	2233-26
2105	2233-30
2106	2233-31
2107	2233-34
2108	2233-36
2109	2233-37

	SEQ ID NO:	Name
	2110	2233-43
	2111	2233-44
	2112	2233-49
5	2113	2233-50
	2114	2233-52
	2115	2233-62
	2116	2233-73
	2117	2233-76
10	2118	2233-78
	2119	2233-79
	2120	2233-82
	2121	2233-88
	2122	2234-04
	2123	2234-13
15	2124	2234-16
	2125	2234-22
	2126	2234-24
	2127	2234-38
	2128	2234-41
20	2129	2234-45
	2130	2234-48
	2131	2234-49
	2132	2234-59
	2133	2234-69
25	2134	2234-83
	2135	2234-84
	2136	2234-87
	2137	2234-92
	2138	2234-95
30	2139	2240-15
	2140	2240-16
	2141	2240-25
	2142	2240-27
	2143	2240-30
35	2144	2240-32
	2145	2240-41
	2146	2240-45
	2147	2240-50
	2148	2240-56
40	2149	2240-65
	2150	2240-68
	2151	2240-73
	2152	2240-78
	2153	2240-79

2154	2240-81
2155	2240-84
2156	2240-96
2157	2241-06
2158	2241-20
2159	2241-32
2160	2241-33
2161	2241-34
2162	2241-43
2163	2241-46
2164	2241-48
2165	2241-50
2166	2241-53
2167	2241-67
2168	2241-71
2169	2241-75
2170	2241-77
2171	2241-79
2172	2241-88
2173	2241-96
2174	2243-05
2175	2243-07
2176	2243-16
2177	2243-23
2178	2243-26
2179	2243-34
2180	2243-40
2181	2243-42
2182	2243-58
2183	2243-60
2184	2243-65
2185	2243-70
2186	2243-80
2187	2243-84
2188	2244-04
2189	2244-07
2190	2244-09
2191	2244-10
2192	2244-17
2193	2244-18
2194	2244-20
2195	2244-22
2196	2244-24
2197	2244-28

	SEQ ID NO	Name
	2198	2244-34
	2299	2244-37
	2200	2244-39
5	2201	2244-41
	2202	2244-46
	2203	2244-50
	2204	2244-51
	2205	2244-56
10	2206	2244-60
	2207	2244-67
	2208	2244-70
	2209	2244-72
	2210	2244-73
	2211	2244-79
15	2212	2244-81
	2213	2244-85
	2214	2244-88
	2215	2244-89
	2216	2244-90
20	2217	2244-92
	2218	2244-93
	2219	2244-94
	2220	2244-95
	2221	2244-96
25	2222	2253-03
	2223	2253-09
	2224	2253-10
	2225	2253-15
	2226	2253-17
30	2227	2253-25
	2228	2253-26
	2229	2253-29
	2230	2253-31
	2231	2253-34
35	2232	2253-45
	2233	2253-47
	2234	2253-48
	2235	2253-49
	2236	2253-51
40	2237	2253-57
	2238	2253-64
	2239	2253-69
	2240	2253-71
	2241	2253-72

2242	2253-74
2243	2253-75
2244	2253-76
2245	2253-77
2246	2253-85
2247	2253-91
2248	2253-93
2249	2254-02
2250	2254-03
2251	2254-05
2252	2254-09
2253	2254-11
2254	2254-19
2255	2254-20
2256	2254-22
2257	2254-34
2258	2254-42
2259	2254-44
2260	2254-46
2261	2254-49
2262	2254-50
2263	2254-52
2264	2254-59
2265	2254-61
2266	2254-64
2267	2254-73
2268	2254-75
2269	2254-77
2270	2254-78
2271	2254-81
2272	2254-82
2273	2254-89
2274	2254-90
2275	2254-94
2276	2255-06
2277	2255-09
2278	2255-11
2279	2255-26
2280	2255-27
2281	2255-29
2282	2255-38
2283	2255-44
2284	2255-48
2285	2255-53

	SEQ ID NO.	Name
	2286	2255-65
	2287	2255-74
	2288	2255-78
	2289	2255-81
5	2290	2255-83
	2291	2255-84
	2292	2255-87
	2293	2255-92
	2294	2255-93
10	2295	2256-02
	2296	2256-09
	2297	2256-10
	2298	2256-25
	2299	2256-37
15	2300	2256-51

2301	2256-53
2302	2256-56
2303	2256-60
2304	2256-63
2305	2256-66
2306	2256-68
2307	2256-71
2308	2256-74
2309	2256-81
2310	2256-83
2311	2256-84
2312	2256-88
2313	2256-95

In one embodiment, a gene or other nucleic acid molecule of the present invention can be an allelic variant that includes a similar but not identical sequence to a *C. felis* nucleic acid sequence of Table I and/or Table II or a complement thereof. For example, an allelic variant of a *C. felis* gene including SEQ ID NO:1 is a gene that occurs at  
5 essentially the same locus (or loci) in the genome as the gene including SEQ ID NO:1, but which, due to natural variations caused by, for example, mutation or recombination, has a similar but not identical sequence. Because natural selection typically selects against alterations that affect function, allelic variants (i.e. alleles corresponding to, or of, cited nucleic acid sequences) usually encode proteins having similar activity to that of the  
10 protein encoded by the gene to which they are being compared. Allelic variants of genes or nucleic acid molecules can also comprise alterations in the 5' or 3' untranslated regions of the gene (e.g., in regulatory control regions), or can involve alternative splicing of a nascent transcript, thereby bringing alternative exons into juxtaposition. Allelic variants are well known to those skilled in the art and would be expected to occur naturally within  
15 a given flea such as *C. felis*, since the genome is diploid, and sexual reproduction will result in the reassortment of alleles.

In one embodiment of the present invention, isolated HMT and HNC proteins are encoded by nucleic acid molecules that hybridize under stringent hybridization conditions to genes or other nucleic acid molecules encoding flea HMT and HNC proteins,  
20 respectively. The minimal size of HMT and HNC proteins of the present invention is a size sufficient to be encoded by a nucleic acid molecule capable of forming a stable hybrid (i.e., hybridizing under stringent hybridization conditions) with the complementary sequence of a nucleic acid molecule encoding the corresponding natural protein. The size of a nucleic acid molecule encoding such a protein is dependent on the  
25 nucleic acid composition and the percent homology between the flea HMT or HNC nucleic acid molecule and the complementary nucleic acid sequence. It can easily be

understood that the extent of homology required to form a stable hybrid under stringent conditions can vary depending on whether the homologous sequences are interspersed throughout a given nucleic acid molecule or are clustered (i.e., localized) in distinct regions on a given nucleic acid molecule.

5           The minimal size of a nucleic acid molecule capable of forming a stable hybrid with a gene encoding a flea HMT or HNC protein is typically at least about 12 to about 15 nucleotides in length if the nucleic acid molecule is GC-rich and at least about 15 to about 17 bases in length if it is AT-rich. The minimal size of a nucleic acid molecule used to encode an HMT or HNC protein homologue of the present invention is from  
10   about 12 to about 18 nucleotides in length. Thus, the minimal size of HMT or HNC protein homologues of the present invention is from about 4 to about 6 amino acids in length. There is no limit, other than a practical limit, on the maximal size of a nucleic acid molecule encoding a flea HMT or HNC protein of the present invention because a nucleic acid molecule of the present invention can include a portion of a gene, an entire  
15   gene, or multiple genes. The preferred size of a protein encoded by a nucleic acid molecule of the present invention depends on whether a full-length, fusion, multivalent, or functional portion of such a protein is desired.

Stringent hybridization conditions are determined based on defined physical properties of the gene to which the nucleic acid molecule is being hybridized, and can be  
20   defined mathematically. Stringent hybridization conditions are those experimental parameters that allow an individual skilled in the art to identify significant similarities between heterologous nucleic acid molecules. These conditions are well known to those skilled in the art. See, for example, Sambrook, *et al.*, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Labs Press, and Meinkoth, *et al.*, 1984, *Anal.*  
25   *Biochem.* 138, 267-284, each of which is incorporated by reference herein in its entirety. As explained in detail in the cited references, the determination of hybridization conditions involves the manipulation of a set of variables including the ionic strength (M,

in moles/liter), the hybridization temperature (°C), the concentration of nucleic acid helix destabilizing agents (such as formamide), the average length of the shortest hybrid duplex (n), and the percent G + C composition of the fragment to which an unknown nucleic acid molecule is being hybridized. For nucleic acid molecules of at least about 150 nucleotides, these variables are inserted into a standard mathematical formula to calculate the melting temperature, or  $T_m$ , of a given nucleic acid molecule. As defined in the formula below,  $T_m$  is the temperature at which two complementary nucleic acid molecule strands will disassociate, assuming 100% complementarity between the two strands:

$$T_m = 81.5^{\circ}\text{C} + 16.6 \log M + 0.41(\%G + C) - 500/n - 0.61(\%\text{formamide}).$$

For nucleic acid molecules smaller than about 50 nucleotides, hybrid stability is defined by the dissociation temperature ( $T_d$ ), which is defined as the temperature at which 50% of the duplexes dissociate. For these smaller molecules, the stability at a standard ionic strength is defined by the following equation:

$$T_d = 4(G + C) + 2(A + T).$$

A temperature of 5°C below  $T_d$  is used to detect hybridization between perfectly matched molecules.

Also well known to those skilled in the art is how base pair mismatch, i.e. differences between two nucleic acid molecules being compared, including non-complementarity of bases at a given location, and gaps due to insertion or deletion of one or more bases at a given location on either of the nucleic acid molecules being compared, will affect  $T_m$  or  $T_d$  for nucleic acid molecules of different sizes. For example,  $T_m$  decreases about 1°C for each 1% of mismatched base pairs for hybrids greater than about 150 bp, and  $T_d$  decreases about 5°C for each mismatched base pair for hybrids below about 50 bp. Conditions for hybrids between about 50 and about 150 base pairs can be determined empirically and without undue experimentation using standard laboratory procedures well known to those skilled in the art. These simple procedures allow one skilled in the art to set the hybridization conditions (by altering, for example, the salt

concentration, the helix destabilizing compound concentration or the temperature) so that only nucleic acid hybrids with greater than a specified % base pair mismatch will hybridize. Stringent hybridization conditions are commonly understood by those skilled in the art to be those experimental conditions that will allow less than or equal to about 5 30% base pair mismatch (i.e., at least about 70% identity). Because one skilled in the art can easily determine whether a given nucleic acid molecule to be tested is less than or greater than about 50 nucleotides, and can therefore choose the appropriate formula for determining hybridization conditions, he or she can determine whether the nucleic acid molecule will hybridize with a given gene under stringent hybridization conditions and 10 similarly whether the nucleic acid molecule will hybridize under conditions designed to allow a desired amount of base pair mismatch.

Hybridization reactions are often carried out by attaching the nucleic acid molecule to be hybridized to a solid support such as a membrane, and then hybridizing with a labeled nucleic acid molecule, typically referred to as a probe, suspended in a 15 hybridization solution. Examples of common hybridization reaction techniques include, but are not limited to, the well-known Southern and northern blotting procedures. Typically, the actual hybridization reaction is done under non-stringent conditions, i.e., at a lower temperature and/or a higher salt concentration, and then high stringency is achieved by washing the membrane in a solution with a higher temperature and/or lower 20 salt concentration in order to achieve the desired stringency.

For example, if the skilled artisan wished to identify a nucleic acid molecule that hybridizes under conditions that would allow less than or equal to 30% pair mismatch with a flea nucleic acid molecule of about 150 bp in length or greater, the following conditions could preferably be used. The average G + C content of flea DNA is about 25 37%, as calculated from known flea nucleic acid sequences. The unknown nucleic acid molecules would be attached to a support membrane, and the 150 bp probe would be labeled, e.g. with a radioactive tag. The hybridization reaction could be carried out in a



solution comprising 2X SSC in the absence of helix destabilizing compounds, at a temperature of about 37°C (low stringency conditions). Solutions of differing concentrations of SSC can be made by one of skill in the art by diluting a stock solution of 20X SSC (175.3 gram NaCl and about 88.2 gram sodium citrate in 1 liter of water, pH 7) to obtain the desired concentration of SSC. The skilled artisan would calculate the washing conditions required to allow up to 30% base pair mismatch. For example, in a wash solution comprising 1X SSC in the absence of helix destabilizing compounds, the  $T_m$  of perfect hybrids would be about 77°C:

$$81.5^{\circ}\text{C} + 16.6 \log (.15\text{M}) + (0.41 \times 0.37) - (500/150) - (0.61 \times 0) = 77.5^{\circ}\text{C}.$$

Thus, to achieve hybridization with nucleic acid molecules having about 30% base pair mismatch, hybridization washes would be carried out at a temperature of less than or equal to 47.5°C. It is thus within the skill of one in the art to calculate additional hybridization temperatures based on the desired percentage base pair mismatch, formulae and G/C content disclosed herein. For example, it is appreciated by one skilled in the art that as the nucleic acid molecule to be tested for hybridization against nucleic acid molecules of the present invention having sequences specified herein becomes longer than 150 nucleotides, the  $T_m$  for a hybridization reaction allowing up to 30% base pair mismatch will not vary significantly from 47.5°C.

Furthermore, it is known in the art that there are commercially available computer programs for determining the degree of similarity between two nucleic acid or protein sequences. These computer programs include various known methods to determine the percentage identity and the number and length of gaps between hybrid nucleic acid molecules or proteins. Preferred methods to determine the percent identity among amino acid sequences and also among nucleic acid sequences include analysis using one or more of the commercially available computer programs designed to compare and analyze nucleic acid or amino acid sequences. These computer programs include, but are not limited to, the SeqLab® Wisconsin Package™ Version 10.0-UNIX sequence analysis

software (hereinafter referred to as "SeqLab"), available from Genetics Computer Group, Madison, WI; and DNAsis® sequence analysis software, version 2.0, available from Hitachi Software, San Bruno, CA. Such software programs represent a collection of algorithms paired with a graphical user interface for using the algorithms. The DNAsis version 2.0 software and SeqLab Wisconsin Package Version 10.0-UNIX software, for example, employ a particular algorithm, the Needleman-Wunsch algorithm to perform pair-wise comparisons between two sequences to yield a percentage identity score, see Needleman, S.B. and Wunsch, C.D., 1970, *J. Mol. Biol.*, 48, 443, which is incorporated herein by reference in its entirety. Such algorithms, including the Needleman-Wunsch algorithm, are commonly used by those skilled in the nucleic acid and amino acid sequencing art to compare sequences. A preferred method to determine percent identity among amino acid sequences and also among nucleic acid sequences includes using the Needleman-Wunsch algorithm, available in the SeqLab Wisconsin Package Version 10.0-UNIX software (hereinafter "SeqLab"), using the Pairwise Comparison/Gap function with the nwsgapdna.cmp scoring matrix, the gap creation penalty and the gap extension penalties set at default values, and the gap shift limits set at maximum (hereinafter referred to as "SeqLab default parameters"). An additional preferred method to determine percent identity among amino acid sequences and also among nucleic acid sequences includes using the Higgins-Sharp algorithm, available in the DNAsis version 2.0 software (hereinafter "DNAsis"), with the gap penalty set at 5, the number of top diagonals set at 5, the fixed gap penalty set at 10, the k-tuple set at 2, the window size set at 5, and the floating gap penalty set at 10. A particularly preferred method to determine percent identity among amino acid sequences and also among nucleic acid sequences includes using the Needleman-Wunsch algorithm available in the SeqLab software, using the SeqLab default parameters.

A preferred flea HMT and/or HNC protein includes a protein encoded by a nucleic acid molecule that hybridizes under conditions that preferably allow less than or

equal to 30% base pair mismatch, preferably under conditions that allow less than or equal to 25% base pair mismatch, preferably under conditions that allow less than or equal to 20% base pair mismatch, preferably under conditions that allow less than or equal to 15% base pair mismatch, preferably under conditions that allow less than or  
5 equal to 10% base pair mismatch and preferably under conditions that allow less than or equal to 5% base pair mismatch with a nucleic acid molecule selected from the group consisting of a nucleic acid sequence complementary to a nucleic acid sequence of Table I and/or Table II.

Another embodiment of the present invention includes a flea HMT and/or HNC  
10 protein encoded by a nucleic acid molecule that hybridizes under conditions comprising, (a) hybridizing in a solution comprising 1X SSC in the absence of helix destabilizing compounds, at a temperature of 37°C and (b) washing in a solution comprising 1X SSC in the absence of helix destabilizing compounds, at a temperature of 47.5°C, to an isolated nucleic acid molecule selected from the group consisting of a nucleic acid  
15 sequence complementary to a nucleic acid sequence of Table I and/or Table II.

Another preferred flea HMT and/or HNC protein of the present invention includes a protein that is encoded by a nucleic acid molecule that is preferably at least 70% identical, preferably at least 75% identical, preferably at least 80% identical, preferably at least 85% identical, preferably at least 90% identical, and preferably at least 95%  
20 identical to a nucleic acid molecule having a nucleic acid sequence of Table I and/or Table II; also preferred are fragments (i.e. portions) of such proteins encoded by nucleic acid molecules that are at least 18 nucleotides. Percent identity as used herein is determined using the Compare function by maximum matching within the program DNAsis Version 2.1 using default parameters.

25 Additional preferred flea HMT and/or HNC proteins of the present invention include proteins having an amino acid sequence encoded by a nucleic acid sequence of Table I and/or Table II, and proteins comprising homologues of a protein encoded by a

nucleic acid sequence of Table I and/or Table II, wherein such a homologue comprises at least one epitope that elicits an immune response against a protein encoded by a nucleic acid sequence of Table I and/or Table II.

In one embodiment, a preferred flea HMT or HNC protein comprises an amino acid sequence of at least 6 amino acids, preferably at least 10 amino acids, preferably at least 15 amino acids, preferably at least 20 amino acids, preferably at least 25 amino acids, preferably at least 30 amino acids, preferably at least 35 amino acids, preferably at least 40 amino acids, preferably at least 50 amino acids, preferably at least 100 amino acids, preferably at least 200 amino acids, preferably at least 250 amino acids, preferably at least 300 amino acids, preferably at least 350 amino acids, and preferably at least 375 amino acids. In another embodiment, preferred flea HMT and HNC proteins comprise full-length proteins, i.e., proteins encoded by full-length coding regions, or post-translationally modified proteins thereof, such as mature proteins from which initiating methionine and/or signal sequences or “pro” sequences have been removed.

A fragment of an HMT and/or HNC protein of the present invention preferably comprises at least 5 amino acids, preferably at least 10 amino acids, preferably at least 15 amino acids, preferably at least 20 amino acids, preferably at least 25 amino acids, preferably at least 30 amino acids, preferably at least 35 amino acids, preferably at least 40 amino acids, preferably at least 45 amino acids, preferably at least 50 amino acids, preferably at least 55 amino acids, preferably at least 60 amino acids, preferably at least 65 amino acids, preferably at least 70 amino acids, preferably at least 75 amino acids, preferably at least 80 amino acids, preferably at least 85 amino acids, preferably at least 90 amino acids, preferably at least 95 amino acids, and preferably at least 100 amino acids in length.

In another embodiment, a preferred flea HMT and/or HNC protein of the present invention is encoded by a nucleic acid molecule comprising at least 15 nucleotides, preferably at least 18 nucleotides, preferably at least 20 nucleotides, preferably at least 25

nucleotides, preferably at least 30 nucleotides, preferably at least 40 nucleotides, preferably at least 50 nucleotides, preferably at least 100 nucleotides, preferably at least 150 nucleotides, preferably at least 350 nucleotides, preferably at least 450 nucleotides, preferably at least 550 nucleotides, preferably at least 650 nucleotides, preferably at least 750 nucleotides, preferably at least 1000 nucleotides, and preferably at least 1100 nucleotides in length. In another embodiment, preferred flea HMT and HNC proteins of the present invention are encoded by nucleic acid molecules comprising apparently full-length HMT or HNC coding regions respectively, i.e., nucleic acid molecules encoding an apparently full-length HMT or HNC proteins.

Preferred flea HMT and HNC proteins of the present invention can be used to develop inhibitors that, when administered to an animal in an effective manner, are capable of protecting that animal from flea infestation. In accordance with the present invention, the ability of an inhibitor of the present invention to protect an animal from flea infestation refers to the ability of that protein to, for example, treat, ameliorate and/or prevent infestation caused by fleas. In particular, the phrase "to protect an animal from flea infestation" refers to reducing the potential for flea population expansion on and around the animal (i.e., reducing the flea burden). Preferably, the flea population size is decreased, optimally to an extent that the animal is no longer bothered by fleas. A host animal, as used herein, is an animal from which fleas can feed by attaching to and feeding through the skin of the animal. Fleas, and other ectoparasites, can live on a host animal for an extended period of time or can attach temporarily to an animal in order to feed. At any given time, a certain percentage of a flea population can be on a host animal whereas the remainder can be in the environment of the animal. Such an environment can include not only adult fleas, but also flea eggs and/or flea larvae. The environment can be of any size such that fleas in the environment are able to jump onto and off of a host animal. For example, the environment of an animal can include plants, such as crops, from which

fleas infest an animal. As such, it is desirable not only to reduce the flea burden on an animal per se, but also to reduce the flea burden in the environment of the animal.

Suitable fleas to target include any flea that is essentially incapable of causing disease in an animal administered an inhibitor of the present invention. As such, fleas to  
 5 target include any flea that produces a protein that can be targeted by an inhibitory compound that inhibits a flea HMT or HNC protein function, thereby resulting in the decreased ability of the parasite to cause disease in an animal. Preferred fleas to target include fleas of the following genera: *Ctenocephalides*, *Cyopsyllus*, *Diamanus* (*Oropsylla*), *Echidnophaga*, *Nosopsyllus*, *Pulex*, *Tunga*, and *Xenopsylla*, with those of the  
 10 species *Ctenocephalides canis*, *Ctenocephalides felis*, *Diamanus montanus*, *Echidnophaga gallinacea*, *Nosopsyllus faciatus*, *Pulex irritans*, *Pulex simulans*, *Tunga penetrans* and *Xenopsylla cheopis* being more preferred, with *C. felis* being preferred. Such fleas are also preferred for the isolation of proteins or nucleic acid molecules of the present invention.

15 One embodiment of a flea HMT and/or HNC protein of the present invention is a fusion protein that includes a flea HMT and/or HNC protein-containing domain attached to one or more fusion segments. Suitable fusion segments for use with the present invention include, but are not limited to, segments that can: enhance a protein's stability; act as an immunopotentiator to enhance an immune response against a flea HMT and/or  
 20 HNC protein; and/or assist in purification of a flea HMT and/or HNC protein (e.g., by affinity chromatography). A suitable fusion segment can be a domain of any size that has the desired function (e.g., imparts increased stability, imparts increased immunogenicity to a protein, and/or simplifies purification of a protein). Fusion segments can be joined to amino and/or carboxyl termini of the flea HMT-containing and/or HNC-containing  
 25 domain of the protein and can be susceptible to cleavage in order to enable straight-forward recovery of a flea HMT and/or HNC protein. Fusion proteins are preferably produced by culturing a recombinant cell transformed with a fusion nucleic acid molecule

that encodes a protein including the fusion segment attached to either the carboxyl and/or amino terminal end of an HMT-containing and/or HNC-containing domain. Preferred fusion segments include a metal binding domain (e.g., a poly-histidine segment); an immunoglobulin binding domain (e.g., Protein A; Protein G; T cell; B cell; Fc receptor or complement protein antibody-binding domains); a sugar binding domain (e.g., a maltose binding domain); and/or a "tag" domain (e.g., at least a portion of  $\beta$ -galactosidase, a strep tag peptide, a T7 tag peptide, a Flag™ peptide, or other domains that can be purified using compounds that bind to the domain, such as monoclonal antibodies). More preferred fusion segments include metal binding domains, such as a poly-histidine segment; a maltose binding domain; a strep tag peptide, such as that available from Biometra in Tampa, FL; and an S10 peptide.

The present invention also includes mimetopes of flea HMT and/or HNC proteins of the present invention. As used herein, a mimetope of a flea HMT and/or HNC protein of the present invention refers to any compound that is able to mimic the activity of such an HMT and/or HNC protein, often because the mimetope has a structure that mimics the particular HMT and/or HNC protein. Mimetopes can be, but are not limited to: peptides that have been modified to decrease their susceptibility to degradation such as all-D retro peptides; anti-idiotypic and/or catalytic antibodies, or fragments thereof; non-proteinaceous immunogenic portions of an isolated protein (e.g., carbohydrate structures); and synthetic or natural organic molecules, including nucleic acids. Such mimetopes can be designed using computer-generated structures of proteins of the present invention. Mimetopes can also be obtained by generating random samples of molecules, such as oligonucleotides, peptides or other organic molecules, and screening such samples by affinity chromatography techniques using the corresponding binding partner.

Another embodiment of the present invention is an isolated nucleic acid molecule comprising a flea HMT and/or HNC nucleic acid molecule, i.e. a nucleic acid molecule

that can be isolated from a HMT cDNA library, from a HNC cDNA library, or from both libraries. As used herein, HMT and HNC nucleic acid molecules has the same meaning as HMT and/or HNC nucleic acid molecule. The identifying characteristics of such nucleic acid molecules are heretofore described. A nucleic acid molecule of the present invention can include an isolated natural flea HMT and/or HNC gene or a homologue thereof, the latter of which is described in more detail below. A nucleic acid molecule of the present invention can include one or more regulatory regions, full-length or partial coding regions, or combinations thereof. The minimal size of a nucleic acid molecule of the present invention is a size sufficient to allow the formation of a stable hybrid (i.e., hybridization under stringent hybridization conditions) with the complementary sequence of another nucleic acid molecule. As such, the minimal size of a HMT and/or HNC nucleic acid molecule of the present invention is from 12 to 18 nucleotides in length. Suitable and preferred fleas from which to isolate nucleic acid molecules of the present invention are disclosed herein. Particularly preferred HMT and/or HNC nucleic acid molecules include *C. felis* HMT and/or HNC nucleic acid molecules.

In accordance with the present invention, an isolated nucleic acid molecule is a nucleic acid molecule that has been removed from its natural milieu (i.e., that has been subjected to human manipulation) and can include DNA, RNA, or derivatives of either DNA or RNA. As such, "isolated" does not reflect the extent to which the nucleic acid molecule has been purified. Isolated flea HMT and/or HNC nucleic acid molecules of the present invention, or homologues thereof, can be isolated from a natural source or produced using recombinant DNA technology (e.g., polymerase chain reaction (PCR) amplification or cloning) or chemical synthesis. Isolated flea HMT and/or HNC nucleic acid molecules, and homologues thereof, can include, for example, natural allelic variants and nucleic acid molecules modified by nucleotide insertions, deletions, substitutions, and/or inversions in a manner such that the modifications do not substantially interfere



with the nucleic acid molecule's ability to encode a HMT and/or HNC protein of the present invention.

A flea HMT and/or HNC nucleic acid molecule homologue can be produced using a number of methods known to those skilled in the art, see, for example, Sambrook et al., *ibid.*, is incorporated by reference herein in its entirety. For example, nucleic acid molecules can be modified using a variety of techniques including, but not limited to, classic mutagenesis and recombinant DNA techniques such as site-directed mutagenesis, chemical treatment, restriction enzyme cleavage, ligation of nucleic acid fragments, PCR amplification, synthesis of oligonucleotide mixtures and ligation of mixture groups to "build" a mixture of nucleic acid molecules, and combinations thereof. Nucleic acid molecule homologues can be selected by hybridization with flea HMT and/or HNC nucleic acid molecules or by screening the function of a protein encoded by the nucleic acid molecule (e.g., ability to elicit an immune response against at least one epitope of a flea HMT or HNC protein or to effect HMT or HNC activity).

An isolated nucleic acid molecule of the present invention can include a nucleic acid sequence that encodes at least one flea HMT or HNC protein of the present invention, examples of such proteins being disclosed herein. Although the phrase "nucleic acid molecule" primarily refers to the physical nucleic acid molecule and the phrase "nucleic acid sequence" primarily refers to the sequence of nucleotides on the nucleic acid molecule, the two phrases can be used interchangeably, especially with respect to a nucleic acid molecule, or a nucleic acid sequence, being capable of encoding a flea HMT or HNC protein.

A preferred nucleic acid molecule of the present invention, when administered to an animal, is capable of protecting that animal from flea infestation. As will be disclosed in more detail below, such a nucleic acid molecule can be, or encode, an antisense RNA, a molecule capable of triple helix formation, a ribozyme, or other nucleic acid-based drug compound. In additional embodiments, a nucleic acid molecule of the present invention

can encode a protective protein (e.g., an HMT or HNC protein of the present invention), the nucleic acid molecule being delivered to the animal, for example, by direct injection (i.e., as a genetic vaccine) or in a vehicle such as a recombinant virus vaccine or a recombinant cell vaccine.

5           In one embodiment of the present invention, a preferred flea HMT and/or HNC nucleic acid molecule includes an isolated nucleic acid molecule that hybridizes under conditions that preferably allow less than or equal to 30% base pair mismatch, preferably under conditions that allow less than or equal to 25% base pair mismatch, preferably under conditions that allow less than or equal to 20% base pair mismatch, preferably  
10 under conditions that allow less than or equal to 15% base pair mismatch, preferably under conditions that allow less than or equal to 10% base pair mismatch and preferably under conditions that allow less than or equal to 5% base pair mismatch with a nucleic acid molecule of Table I and/or Table II and/or a nucleic acid molecule that is complementary to a nucleic acid molecule of Table I and/or Table II.

15           Another embodiment of the present invention includes a HMT and/or HNC nucleic acid molecule, wherein said nucleic acid molecule hybridizes, in a solution comprising 1X SSC in the absence of helix destabilizing compounds, at a temperature of 47.5°C, to an isolated nucleic acid molecule a nucleic acid molecule of Table I or Table II and/or a nucleic acid molecule that is complementary to a nucleic acid molecule of Table  
20 I or Table II. Additional preferred nucleic acid molecules of the present invention include oligonucleotides of an isolated nucleic acid molecule, wherein said nucleic acid molecule hybridizes, in a solution comprising 1X SSC in the absence of helix destabilizing compounds, at a temperature of 47.5°C, to an isolated nucleic acid molecule of Table I or Table II and/or a nucleic acid molecule that is complementary to a  
25 nucleic acid molecule of Table I or Table II, wherein said oligonucleotide comprises at least 18 nucleotides.

Additional preferred flea HMT and/or HNC nucleic acid molecules of the present invention include nucleic acid molecules comprising a nucleic acid sequence that is preferably at least 70%, preferably at least 75%, preferably at least 80% preferably at least 85%, preferably at least 90%, and preferably at least 95% identical to a nucleic acid sequence of Table I or Table II and/or a nucleic acid molecule that is complementary to a nucleic acid molecule of Table I or Table II. Also preferred are oligonucleotides of any of such nucleic acid molecules. Percent identity may be determined using the SeqLab sequence analysis software, using default parameters.

Another preferred nucleic acid molecule of the present invention includes at least a portion of nucleic acid sequence of Table I or Table II, as well as allelic variants of nucleic acid molecules having these nucleic acid sequences and homologues of nucleic acid molecules having these nucleic acid sequences; preferably such a homologue encodes or is complementary to a nucleic acid molecule that encodes at least one epitope that elicits an immune response against a protein encoded by a nucleic acid molecule of Table I and/or Table II. Such nucleic acid molecules can include nucleotides in addition to those included in the SEQ ID NOs, such as, but not limited to, a full-length gene, a full-length coding region, a nucleic acid molecule encoding a fusion protein, or a nucleic acid molecule encoding a multivalent protective compound.

In one embodiment, HMT and/or HNC nucleic acid molecule of the present invention encodes a protein that is at least 70%, preferably at least 75%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 95%, preferably at least 98%, and preferably at least 100% identical to a protein encoded by a nucleic acid molecule having a sequence of Table I and/or Table II.

In one embodiment, a HMT and/or HNC nucleic acid molecule of the present invention encodes a protein having an amino acid sequence that is at least 70%, preferably at least 75%, preferably at least 80%, preferably at least 85%, preferably at least 90%, preferably at least 95%, preferably at least 98%, and preferably at least

100% identical to a protein encoded by a nucleic acid molecule having a sequence of Table I and/or Table II. The present invention also includes a HMT and/or HNC nucleic acid molecule encoding a protein having at least a portion of a protein encoded by a nucleic acid molecule having a sequence of Table I and/or Table II, as well as allelic  
 5 variants of a nucleic acid molecule encoding a protein having these sequences, including nucleic acid molecules that have been modified to accommodate codon usage properties of the cells in which such nucleic acid molecules are to be expressed.

In another embodiment, a preferred flea HMT and/or HNC nucleic acid molecule of the present invention comprises a nucleic acid molecule comprising at least 15  
 10 nucleotides, preferably at least 18 nucleotides, preferably at least 20 nucleotides, preferably at least 25 nucleotides, preferably at least 30 nucleotides, preferably at least 40 nucleotides, preferably at least 50 nucleotides, preferably at least 100 nucleotides, preferably at least 150 nucleotides, preferably at least 350 nucleotides, preferably at least 450 nucleotides, preferably at least 550 nucleotides, preferably at least 650 nucleotides,  
 15 preferably at least 750 nucleotides, preferably at least 1000 nucleotides, and preferably at least 1100 nucleotides in length.

In another embodiment, a preferred flea HMT and/or HNC nucleic acid molecule encodes a protein comprising at least 5 amino acids, preferably at least 6 amino acids, preferably at least 10 amino acids, preferably at least 15 amino acids, preferably at least  
 20 20 amino acids, preferably at least 25 amino acids, preferably at least 30 amino acids, preferably at least 40 amino acids, preferably at least 50 amino acids, preferably at least 100 amino acids, preferably at least 150 amino acids, preferably at least 200 amino acids, preferably at least 300 amino acids, preferably at least 350 amino acids, and preferably at least 350 amino acids in length.

25 In another embodiment, a preferred flea HMT and/or HNC nucleic acid molecule of the present invention comprises an apparently full-length HMT and/or HNC coding region, i.e., the preferred nucleic acid molecule encodes an apparently full-length HMT

and/or HNC protein, or a post-translationally modified protein thereof. In one embodiment, a preferred HMT and/or HNC nucleic acid molecule of the present invention encodes a mature protein.

Knowing the nucleic acid sequences of certain flea HMT and/or HNC nucleic acid molecules of the present invention allows one skilled in the art to, for example, (a) make copies of those nucleic acid molecules, (b) obtain nucleic acid molecules including at least a portion of such nucleic acid molecules (e.g., nucleic acid molecules including full-length genes, full-length coding regions, regulatory control sequences, truncated coding regions), and (c) obtain other flea HMT and/or HNC nucleic acid molecules. Such nucleic acid molecules can be obtained in a variety of ways including screening appropriate expression libraries with antibodies of the present invention; traditional cloning techniques using oligonucleotide probes of the present invention to screen appropriate libraries; and PCR amplification of appropriate libraries or DNA using oligonucleotide primers of the present invention. Preferred libraries to screen or from which to amplify nucleic acid molecules include flea 1<sup>st</sup> instar larvae; 3<sup>rd</sup> instar larvae, wandering larvae, prepupal larvae, pupae and whole adult flea cDNA libraries as well as genomic DNA libraries. Similarly, preferred DNA sources to screen or from which to amplify nucleic acid molecules include flea prepupal cDNA, adult cDNA and genomic DNA. Techniques to clone and amplify genes are disclosed, for example, in Sambrook et al., *ibid*.

The present invention also includes nucleic acid molecules that are oligonucleotides capable of hybridizing, under stringent hybridization conditions, with complementary regions of other, preferably longer, nucleic acid molecules of the present invention such as those comprising *C. felis* HMT and/or HNC nucleic acid molecules or other flea HMT and/or HNC nucleic acid molecules. Oligonucleotides of the present invention can be RNA, DNA, or derivatives of either. The minimum size of such oligonucleotides is the size required for formation of a stable hybrid between an

oligonucleotide and a complementary sequence on a nucleic acid molecule of the present invention. A preferred oligonucleotide of the present invention has a maximum size of preferably 100 to 200 nucleotides. The present invention includes oligonucleotides that can be used as, for example, probes to identify nucleic acid molecules, primers to  
5 produce nucleic acid molecules, or therapeutic reagents to inhibit flea HMT and/or HNC protein production or activity (e.g., as antisense, triplex formation, ribozyme and/or RNA drug-based reagents). The present invention also includes the use of such oligonucleotides to protect animals from disease using one or more of such technologies. Appropriate oligonucleotide-containing therapeutic compositions can be administered to  
10 an animal using techniques known to those skilled in the art.

One embodiment of the present invention includes a recombinant vector, which includes at least one isolated nucleic acid molecule of the present invention, inserted into any vector capable of delivering the nucleic acid molecule into a host cell. Such a vector contains heterologous nucleic acid sequences, that is nucleic acid sequences that are not  
15 naturally found adjacent to nucleic acid molecules of the present invention and that preferably are derived from a species other than the species from which the nucleic acid molecule(s) are derived. The vector can be either RNA or DNA, either prokaryotic or eukaryotic, and typically is a virus or a plasmid. Recombinant vectors can be used in the cloning, sequencing, and/or otherwise manipulating of flea HMT and/or HNC nucleic  
20 acid molecules of the present invention.

One type of recombinant vector, referred to herein as a recombinant molecule, comprises a nucleic acid molecule of the present invention operatively linked to an expression vector. The phrase operatively linked refers to insertion of a nucleic acid molecule into an expression vector in a manner such that the molecule is able to be  
25 expressed when transformed into a host cell. As used herein, an expression vector is a DNA or RNA vector that is capable of transforming a host cell and of effecting expression of a specified nucleic acid molecule. Preferably, the expression vector is also

capable of replicating within the host cell. Expression vectors can be either prokaryotic or eukaryotic, and are typically viruses or plasmids. Expression vectors of the present invention include any vectors that function (i.e., direct gene expression) in recombinant cells of the present invention, including in bacterial, fungal, parasite, insect, other animal, and plant cells. Preferred expression vectors of the present invention can direct gene expression in bacterial, yeast, insect and mammalian cells, and preferably in the cell types disclosed herein.

In particular, expression vectors of the present invention contain regulatory sequences such as transcription control sequences, translation control sequences, origins of replication, and other regulatory sequences that are compatible with the recombinant cell and that control the expression of nucleic acid molecules of the present invention. In particular, recombinant molecules of the present invention include transcription control sequences. Transcription control sequences are sequences that control the initiation, elongation, and termination of transcription. Particularly important transcription control sequences are those which control transcription initiation, such as promoter, enhancer, operator and repressor sequences. Suitable transcription control sequences include any transcription control sequence that can function in at least one of the recombinant cells of the present invention. A variety of such transcription control sequences are known to those skilled in the art. Preferred transcription control sequences include those that function in bacterial, yeast, or insect and mammalian cells, such as, but not limited to, *tac*, *lac*, *trp*, *trc*, oxy-pro, omp/lpp, rrnB, bacteriophage lambda (such as lambda p<sub>L</sub> and lambda p<sub>R</sub> and fusions that include such promoters), bacteriophage T7, T7*lac*, bacteriophage T3, bacteriophage SP6, bacteriophage SP01, metallothionein, alpha-mating factor, *Pichia* alcohol oxidase, alphavirus subgenomic promoter, antibiotic resistance gene, baculovirus, *Heliothis zea* insect virus, vaccinia virus, herpesvirus, raccoon poxvirus, other poxvirus, adenovirus, cytomegalovirus (such as immediate early promoter), simian virus 40, retrovirus, actin, retroviral long terminal repeat, Rous

sarcoma virus, heat shock, phosphate and nitrate transcription control sequences as well as other sequences capable of controlling gene expression in prokaryotic or eukaryotic cells. Additional suitable transcription control sequences include tissue-specific promoters and enhancers as well as lymphokine-inducible promoters (e.g., promoters  
5 inducible by interferons or interleukins). Transcription control sequences of the present invention can also include naturally occurring transcription control sequences naturally associated with fleas, such as *C. felis* transcription control sequences.

Suitable and preferred nucleic acid molecules to include in recombinant vectors of the present invention are as disclosed herein. Preferred nucleic acid molecules to include  
10 in recombinant vectors, and particularly in recombinant molecules, include nucleic acid molecules having a sequence of Table I and/or Table II.

Recombinant molecules of the present invention may also (a) contain secretory signals (i.e., signal segment nucleic acid sequences) to enable an expressed flea protein of the present invention to be secreted from the cell that produces the protein and/or (b)  
15 contain fusion sequences which lead to the expression of nucleic acid molecules of the present invention as fusion proteins. Examples of suitable signal segments include any signal segment capable of directing the secretion of a protein of the present invention. Preferred signal segments include, but are not limited to, tissue plasminogen activator (t-PA), interferon, interleukin, growth hormone, histocompatibility and viral envelope  
20 glycoprotein signal segments. Suitable fusion segments encoded by fusion segment nucleic acids are disclosed herein. In addition, a nucleic acid molecule of the present invention can be joined to a fusion segment that directs the encoded protein to the proteosome, such as a ubiquitin fusion segment. Eukaryotic recombinant molecules may also include intervening and/or untranslated sequences surrounding and/or within the  
25 nucleic acid sequences of nucleic acid molecules of the present invention.

Another embodiment of the present invention includes a recombinant cell comprising a host cell transformed with one or more recombinant molecules of the



present invention. Transformation of a nucleic acid molecule into a cell can be accomplished by any method by which a nucleic acid molecule can be inserted into the cell. Transformation techniques include, but are not limited to, transfection, electroporation, microinjection, lipofection, adsorption, and protoplast fusion. A  
5 recombinant cell may remain unicellular or may grow into a tissue, organ or a multicellular organism. It is to be noted that a cell line refers to any recombinant cell of the present invention that is not a transgenic animal. Transformed nucleic acid molecules of the present invention can remain extrachromosomal or can integrate into one or more sites within a chromosome of the transformed (i.e., recombinant) cell in such a manner  
10 that their ability to be expressed is retained. Preferred nucleic acid molecules with which to transform a cell include *C. felis* HMT and HNC nucleic acid molecules disclosed herein. Preferred nucleic acid molecules with which to transform a cell include nucleic acid molecules having a sequence of Table I and/or Table II.

Suitable host cells to transform include any cell that can be transformed with a  
15 nucleic acid molecule of the present invention. Host cells can be either untransformed cells or cells that are already transformed with at least one nucleic acid molecule (e.g., nucleic acid molecules encoding one or more proteins of the present invention and/or other proteins useful in the production of multivalent vaccines). Host cells of the present invention either can be endogenously (i.e., naturally) capable of producing flea HMT  
20 and/or HNC proteins of the present invention or can be capable of producing such proteins after being transformed with at least one nucleic acid molecule of the present invention. Host cells of the present invention can be any cell capable of producing at least one protein of the present invention, and include bacterial, fungal (including yeast), parasite (including helminth, protozoa and ectoparasite), other insect, other animal and  
25 plant cells. Preferred host cells include bacterial, mycobacterial, yeast, insect and mammalian cells. More preferred host cells include *Salmonella*, *Escherichia*, *Bacillus*, *Caulobacter*, *Listeria*, *Saccharomyces*, *Pichia*, *Spodoptera*, *Mycobacteria*, *Trichoplusia*,

BHK (baby hamster kidney) cells, MDCK cells (Madin-Darby canine kidney cell line), CRFK cells (Crandell feline kidney cell line), CV-1 cells (African monkey kidney cell line used, for example, to culture raccoon poxvirus), COS (e.g., COS-7) cells, and Vero cells. Particularly preferred host cells are *Escherichia coli*, including *E. coli* K-12 derivatives; *Salmonella typhi*; *Salmonella typhimurium*, including attenuated strains such as UK-1  $\chi$ 3987 and SR-11  $\chi$ 4072; *Caulobacter*; *Pichia*; *Spodoptera frugiperda*; *Trichoplusia ni*; BHK cells; MDCK cells; CRFK cells; CV-1 cells; COS cells; Vero cells; and non-tumorigenic mouse myoblast G8 cells (e.g., ATCC CRL 1246). Additional appropriate mammalian cell hosts include other kidney cell lines, other fibroblast cell lines (e.g., human, murine or chicken embryo fibroblast cell lines), myeloma cell lines, Chinese hamster ovary cells, mouse NIH/3T3 cells, LMTK<sup>31</sup> cells and/or HeLa cells. In one embodiment, the proteins may be expressed as heterologous proteins in myeloma cell lines employing immunoglobulin promoters.

A recombinant cell is preferably produced by transforming a host cell with one or more recombinant molecules, each comprising one or more nucleic acid molecules of the present invention operatively linked to an expression vector containing one or more transcription control sequences, examples of which are disclosed herein. The phrase operatively linked refers to insertion of a nucleic acid molecule into an expression vector in a manner such that the molecule is able to be expressed when transformed into a host cell.

A recombinant cell of the present invention includes any cell transformed with at least one of any nucleic acid molecule of the present invention. Suitable and preferred nucleic acid molecules as well as suitable and preferred recombinant molecules with which to transform cells are disclosed herein.

Recombinant cells of the present invention can also be co-transformed with one or more recombinant molecules including flea HMT and/or HNC nucleic acid molecules encoding one or more proteins of the present invention and one or more other nucleic

acid molecules encoding other protective compounds, as disclosed herein (e.g., to produce multivalent vaccines).

Recombinant DNA technologies can be used to improve expression of transformed nucleic acid molecules by manipulating, for example, the number of copies of the nucleic acid molecules within a host cell, the efficiency with which those nucleic acid molecules are transcribed, the efficiency with which the resultant transcripts are translated, and the efficiency of post-translational modifications. Recombinant techniques useful for increasing the expression of nucleic acid molecules of the present invention include, but are not limited to, operatively linking nucleic acid molecules to high-copy number plasmids, integration of the nucleic acid molecules into one or more host cell chromosomes, addition of vector stability sequences to plasmids, substitutions or modifications of transcription control signals (e.g., promoters, operators, enhancers), substitutions or modifications of translational control signals (e.g., ribosome binding sites, Shine-Dalgarno sequences), modification of nucleic acid molecules of the present invention to correspond to the codon usage of the host cell, deletion of sequences that destabilize transcripts, and use of control signals that temporally separate recombinant cell growth from recombinant enzyme production during fermentation. The activity of an expressed recombinant protein of the present invention may be improved by fragmenting, modifying, or derivatizing nucleic acid molecules encoding such a protein.

Isolated flea HMT and/or HNC proteins of the present invention can be produced in a variety of ways, including production and recovery of natural proteins, production and recovery of recombinant proteins, and chemical synthesis of the proteins. In one embodiment, an isolated protein of the present invention is produced by culturing a cell capable of expressing the protein under conditions effective to produce the protein, and recovering the protein. A preferred cell to culture is a recombinant cell of the present invention. Effective culture conditions include, but are not limited to, effective media, bioreactor, temperature, pH and oxygen conditions that permit protein production. An

effective, medium refers to any medium in which a cell is cultured to produce a flea HMT and/or HNC protein of the present invention. Such medium typically comprises an aqueous medium having assimilable carbon, nitrogen and phosphate sources, and appropriate salts, minerals, metals and other nutrients, such as vitamins. Cells of the present invention can be cultured in conventional fermentation bioreactors, shake flasks, test tubes, microtiter dishes, and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a recombinant cell. Such culturing conditions are within the expertise of one of ordinary skill in the art. Examples of suitable conditions are included in the Examples section.

Depending on the vector and host system used for production, resultant proteins of the present invention may either remain within the recombinant cell; be secreted into the fermentation medium; be secreted into a space between two cellular membranes, such as the periplasmic space in *E. coli*; or be retained on the outer surface of a cell or viral membrane.

The phrase "recovering the protein", as well as similar phrases, refers to collecting the whole fermentation medium containing the protein and need not imply additional steps of separation or purification. Proteins of the present invention can be purified using a variety of standard protein purification techniques, such as, but not limited to, affinity chromatography, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential solubilization. Proteins of the present invention are preferably retrieved in "substantially pure" form. As used herein, "substantially pure" refers to a purity that allows for the effective use of the protein as a therapeutic composition or diagnostic. A therapeutic composition for animals, for example, should exhibit no substantial toxicity and preferably should be capable of stimulating the production of antibodies in a treated animal.

The present invention also includes isolated (i.e., removed from their natural milieu) antibodies that selectively bind to a flea HMT and/or HNC protein of the present invention or a mimetope thereof (e.g., anti-*C. felis* HMT or HNC antibodies). As used herein, the term "selectively binds to" an HMT and/or HNC protein refers to the ability of antibodies of the present invention to preferentially bind to specified proteins and mimetopes thereof of the present invention. Binding can be measured using a variety of methods standard in the art including enzyme immunoassays (e.g., ELISA), immunoblot assays, etc.; see, for example, Sambrook et al., *ibid.*, and Harlow, et al., 1988, *Antibodies, a Laboratory Manual*, Cold Spring Harbor Labs Press; Harlow et al., *ibid.*, is incorporated by reference herein in its entirety. An anti-HMT or anti-HNC antibody of the present invention preferably selectively binds to a flea HMT or HNC protein respectively in such a way as to inhibit the function of that protein.

Isolated antibodies of the present invention can include antibodies in serum, or antibodies that have been purified to varying degrees. Antibodies of the present invention can be polyclonal or monoclonal, or can be functional equivalents such as antibody fragments and genetically-engineered antibodies, including single chain antibodies or chimeric antibodies that can bind to one or more epitopes.

A preferred method to produce antibodies of the present invention includes (a) administering to an animal an effective amount of a protein, peptide or mimetope thereof of the present invention to produce the antibodies and (b) recovering the antibodies. In another method, antibodies of the present invention are produced recombinantly using techniques as heretofore disclosed to produce HMT and/or HNC proteins of the present invention. Antibodies raised against defined proteins or mimetopes can be advantageous because such antibodies are not substantially contaminated with antibodies against other substances that might otherwise cause interference in a diagnostic assay or side effects if used in a therapeutic composition.

Antibodies of the present invention have a variety of potential uses that are within the scope of the present invention. For example, such antibodies can be used (a) as therapeutic compounds to passively immunize an animal in order to protect the animal from fleas susceptible to treatment by such antibodies and/or (b) as tools to screen  
5 expression libraries and/or to recover desired proteins of the present invention from a mixture of proteins and other contaminants. Furthermore, antibodies of the present invention can be used to target cytotoxic agents to fleas in order to directly kill such fleas. Targeting can be accomplished by conjugating (i.e., stably joining) such antibodies to the cytotoxic agents using techniques known to those skilled in the art. Suitable cytotoxic  
10 agents are known to those skilled in the art.

One embodiment of the present invention is a therapeutic composition that, when administered to an animal susceptible to flea infestation, is capable of protecting that animal from flea infestation. Therapeutic compositions of the present invention include at least one of the following protective molecules: an isolated flea HMT and/or HNC  
15 protein; a mimetope of an isolated flea HMT and/or HNC protein; an isolated flea HMT and/or HNC nucleic acid molecule; and/or a compound derived from said isolated flea HMT and/or HNC protein that inhibits HMT and/or HNC protein activity. A therapeutic composition of the present invention can further comprise a component selected from the group of an excipient, a carrier, and/or an adjuvant; these components are described  
20 further herein. As used herein, a protective molecule or protective compound refers to a compound that, when administered to an animal in an effective manner, is able to treat, ameliorate, and/or prevent flea infestation. Preferred fleas to target are heretofore disclosed. One example of a protective molecule is a vaccine, such as, but not limited to, a naked nucleic acid vaccine, a recombinant virus vaccine, a recombinant cell vaccine,  
25 and a recombinant protein vaccine. Another example of a protective molecule is a compound that inhibits HMT and/or HNC protein activity, such as an isolated antibody that selectively binds to a flea HMT and/or HNC protein, a substrate analog of a flea

HMT and/or HNC protein, anti-sense-, triplex formation-, ribozyme-, and/or RNA drug-based compounds, or other inorganic or organic molecules that inhibit HMT and/or HNC protein activity. Inhibiting flea HMT and/or HNC protein activity can refer to the ability of a compound to reduce the activity of flea HMT and/or HNC proteins. Inhibiting flea  
5 HMT and/or HNC protein activity can also refer to the ability of a compound to reduce the amount of flea HMT and/or HNC protein in a flea.

Another embodiment of the present invention includes a method to reduce a flea infestation in an animal susceptible to flea infestation. Such a method includes the step of administering to the animal a therapeutic molecule comprising a protective compound  
10 selected from the group consisting of (a) an isolated flea HMT and/or HNC protein; (b) a mimetope of an isolated flea HMT and/or HNC protein; (c) an isolated flea HMT and/or HNC nucleic acid molecule; and (d) a compound derived from an isolated flea HMT and/or HNC protein that inhibits HMT and/or HNC protein activity.

Therapeutic compositions of the present invention can be administered to any  
15 animal susceptible to flea infestation, preferably to mammals, and preferably to dogs, cats, humans, ferrets, horses, cattle, sheep, and other pets, economic food animals, work animals and/or zoo animals. Preferred animals to protect against flea infestation include dogs, cats, humans, and ferrets, with dogs and cats being particularly preferred.

As used herein, the term derived, or the term derived from, refers to a peptide,  
20 antibody, mimetope, nucleic acid molecule, or other compound that was obtained from or obtained using a flea HMT and/or HNC protein or nucleic acid molecule of the present invention. Methods to obtain derivatives from a HMT and/or HNC molecule of the present invention are known in the art, and as such include, but are not limited to molecular modeling of HMT and/or HNC proteins to determine active sites, i.e. sites that  
25 interact with other molecules, and predicting from these active sites smaller fragments and/or mimetopes that retain and/or mimic these active sites, thereby inhibiting HMT and/or HNC protein activity; screening of peptide or small chemical compound libraries

against HMT and/or HNC proteins of the present invention; and screening of polyclonal or monoclonal antibodies to find antibodies that specifically bind HMT and/or HNC proteins of the present invention.

An HMT and/or HNC protein inhibitor of the present invention is identified by its  
5 ability to bind to, modify, or otherwise interact with, a flea HMT and/or HNC protein, thereby inhibiting the activity of HMT and/or HNC proteins. Suitable inhibitors of HMT and/or HNC protein activity are compounds that inhibit HMT and/or HNC protein activity in at least one of a variety of ways: (a) by binding to or otherwise interacting with or otherwise modifying HMT and/or HNC protein sites; (b) by binding to the HMT  
10 and/or HNC protein and thus reducing the availability of the HMT and/or HNC protein in solution; and (c) by interacting with other regions of the HMT and/or HNC protein to inhibit HMT and/or HNC protein activity, for example, by allosteric interaction.

Flea HMT and/or HNC protein inhibitors can be used directly as compounds in compositions of the present invention to treat animals as long as such compounds are not  
15 harmful to host animals being treated. Preferred HMT and/or HNC protein inhibitors of the present invention include, but are not limited to, flea HMT and/or HNC protein substrate analogs, and other molecules that bind to a flea HMT and/or HNC proteins (e.g., to an allosteric site) in such a manner that the activity of the flea HMT and/or HNC protein is inhibited. An HMT and/or HNC protein substrate analog refers to a compound  
20 that interacts with (e.g., binds to, associates with, modifies) the active site of a HMT and/or HNC protein. A preferred HMT and/or HNC protein substrate analog inhibits HMT and/or HNC protein activity. HMT and/or HNC protein substrate analogs can be of any inorganic or organic composition. HMT and/or HNC protein substrate analogs can be, but need not be, structurally similar to a HMT and/or HNC protein natural substrate  
25 as long as they can interact with the active site of that HMT and/or HNC protein. HMT and/or HNC protein substrate analogs can be designed using computer-generated structures of HMT and/or HNC proteins of the present invention or computer structures



of HMT and/or HNC protein's natural substrates. Preferred sites to model include one or more of the active sites of HMT and/or HNC protein. Substrate analogs can also be obtained by generating random samples of molecules, such as oligonucleotides, peptides, peptidomimetic compounds, or other inorganic or organic molecules, and screening such samples for their ability to interfere with interaction between HMT and/or HNC proteins and their substrates, e.g. by affinity chromatography techniques. A preferred HMT and/or HNC protein substrate analog is a HMT and/or HNC protein mimetic compound, i.e., a compound that is structurally and/or functionally similar to a natural substrate of a HMT and/or HNC protein of the present invention, particularly to the region of the substrate that interacts with the HMT and/or HNC protein active site, but that inhibits HMT and/or HNC protein activity upon interacting with the HMT and/or HNC protein active site.

The present invention also includes a therapeutic composition comprising at least one protective molecule of the present invention in combination with at least one additional compound protective against one or more infectious agents.

In one embodiment, a therapeutic composition of the present invention can be used to protect an animal from flea infestation by administering such composition to a flea in order to prevent infestation. Such administration to the flea and/or animal could be oral, or by application to the animal's body surface (e.g. topical spot-on, or spraying onto the animal), or by application to the environment (e.g., spraying). Examples of such compositions include, but are not limited to, transgenic vectors capable of producing at least one therapeutic composition of the present invention. In another embodiment a flea can ingest therapeutic compositions, or products thereof, present on the surface of or in the blood of a host animal that has been administered a therapeutic composition of the present invention.

In accordance with the present invention, a host animal (i.e., an animal that is or is capable of being infested with fleas) is treated by administering to the animal a

therapeutic composition of the present invention in such a manner that the composition itself (e.g., a HMT and/or HNC protein, a HMT and/or HNC nucleic acid molecule, a HMT and/or HNC protein inhibitor, a HMT and/or HNC protein synthesis suppressor (i.e., a compound that decreases the production or half-life of a HMT and/or HNC protein in fleas), a HMT and/or HNC protein mimetope, or a anti-HMT and/or HNC antibody) or a product generated by the animal in response to administration of the composition (e.g., antibodies produced in response to administration of a flea HMT and/or HNC protein or nucleic acid molecule, or conversion of an inactive inhibitor "prodrug" to an active HMT and/or HNC protein inhibitor) ultimately enters the flea. A host animal is preferably treated in such a way that the compound or product thereof is present on the body surface of the animal or enters the blood stream of the animal. Fleas are then exposed to the composition or product when they feed from the animal. For example, flea HMT and/or HNC protein inhibitors administered to an animal are administered in such a way that the inhibitors enter the blood stream of the animal, where they can be taken up by feeding fleas.

The present invention also includes the ability to reduce larval flea infestation in that when fleas feed from a host animal that has been administered a therapeutic composition of the present invention, at least a portion of compounds of the present invention, or products thereof, in the blood taken up by the fleas are excreted by the fleas in feces, which is subsequently ingested by flea larvae. In particular, it is of note that flea larvae obtain most, if not all, of their nutrition from flea feces.

In accordance with the present invention, reducing HMT and/or HNC protein activity in a flea can lead to a number of outcomes that reduce flea burden on treated animals and their surrounding environments. Such outcomes include, but are not limited to, (a) reducing the viability of fleas that feed from the treated animal, (b) reducing the fecundity of female fleas that feed from the treated animal, (c) reducing the reproductive capacity of male fleas that feed from the treated animal, (d) reducing the viability of eggs

laid by female fleas that feed from the treated animal, (e) altering the blood feeding behavior of fleas that feed from the treated animal (e.g., fleas take up less volume per feeding or feed less frequently), (f) reducing the viability of flea larvae, for example due to the feeding of larvae from feces of fleas that feed from the treated animal, (g) altering  
5 the development of flea larvae (e.g., by decreasing feeding behavior, inhibiting growth, inhibiting (e.g., slowing or blocking) molting, and/or otherwise inhibiting maturation to adults), and/or (h) altering or decreasing the ability of fleas or flea larvae to digest a blood meal.

In order to protect an animal from flea infestation, a therapeutic composition of  
10 the present invention is administered to the animal in an effective manner such that the composition is capable of protecting that animal from flea infestation. Therapeutic compositions of the present invention can be administered to animals prior to infestation in order to prevent infestation (i.e., as a preventative vaccine) and/or can be administered to animals after infestation. For example, proteins, mimetopes thereof, and antibodies  
15 thereof can be used as immunotherapeutic agents.

Therapeutic compositions of the present invention can be formulated in an excipient that the animal to be treated can tolerate. Examples of such excipients include water, saline, Ringer's solution, dextrose solution, Hank's solution, and other aqueous physiologically balanced salt solutions. Nonaqueous vehicles, such as fixed oils, sesame  
20 oil, ethyl oleate, or triglycerides may also be used. Other useful formulations include suspensions containing viscosity enhancing agents, such as sodium carboxymethylcellulose, sorbitol, or dextran. Excipients can also contain minor amounts of additives, such as substances that enhance isotonicity and chemical stability. Examples of buffers include phosphate buffer, bicarbonate buffer and Tris buffer, while  
25 examples of preservatives include thimerosal, or o-cresol, formalin and benzyl alcohol. Standard formulations can either be liquid injectables or solids which can be taken up in a suitable liquid as a suspension or solution for injection. Thus, in a non-liquid

formulation, the excipient can comprise dextrose, human serum albumin, preservatives, etc., to which sterile water or saline can be added prior to administration.

In one embodiment of the present invention, a therapeutic composition can include an adjuvant. Adjuvants are agents that are capable of enhancing the immune response of an animal to a specific antigen. Suitable adjuvants include, but are not limited to, cytokines, chemokines, and compounds that induce the production of cytokines and chemokines (e.g., granulocyte macrophage colony stimulating factor (GM-CSF), Flt-3 ligand, granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), colony stimulating factor (CSF), erythropoietin (EPO), interleukin 2 (IL-2), interleukin-3 (IL-3), interleukin 4 (IL-4), interleukin 5 (IL-5), interleukin 6 (IL-6), interleukin 7 (IL-7), interleukin 8 (IL-8), interleukin 10 (IL-10), interleukin 12 (IL-12), interferon gamma, interferon gamma inducing factor I (IGIF), transforming growth factor beta, RANTES (regulated upon activation, normal T cell expressed and presumably secreted), macrophage inflammatory proteins (e.g., MIP-1 alpha and MIP-1 beta), and Leishmania elongation initiating factor (LEIF)); bacterial components (e.g., endotoxins, in particular superantigens, exotoxins and cell wall components); aluminum-based salts; calcium-based salts; silica; polynucleotides; toxoids; serum proteins, viral coat proteins; block copolymer adjuvants (e.g., Hunter's Titermax<sup>TM</sup> adjuvant (Vaxcel<sup>TM</sup>, Inc. Norcross, GA), Ribi adjuvants (Ribi ImmunoChem Research, Inc., Hamilton, MT); and saponins and their derivatives (e.g., Quil A (Superfos Biosector A/S, Denmark). Protein adjuvants of the present invention can be delivered in the form of the protein themselves or of nucleic acid molecules encoding such proteins using the methods described herein.

In one embodiment of the present invention, a therapeutic composition can include a carrier. Carriers include compounds that increase the half-life of a therapeutic composition in the treated animal. Suitable carriers include, but are not limited to,

polymeric controlled release vehicles, biodegradable implants, liposomes, bacteria, viruses, other cells, oils, esters, and glycols.

One embodiment of the present invention is a controlled release formulation that is capable of slowly releasing a composition of the present invention into an animal. As  
5 used herein, a controlled release formulation comprises a composition of the present invention in a controlled release vehicle. Suitable controlled release vehicles include, but are not limited to, biocompatible polymers, other polymeric matrices, capsules, microcapsules, microparticles, bolus preparations, osmotic pumps, diffusion devices, liposomes, lipospheres, and transdermal delivery systems. Other controlled release  
10 formulations of the present invention include liquids that, upon administration to an animal, form a solid or a gel *in situ*. Preferred controlled release formulations are biodegradable (i.e., bioerodible).

A preferred controlled release formulation of the present invention is capable of releasing a composition of the present invention into the blood of the treated animal at a  
15 constant rate sufficient to attain therapeutic dose levels of the composition to protect an animal from flea infestation. The therapeutic composition is preferably released over a period of time ranging from about 1 to about 12 months. A controlled release formulation of the present invention is capable of effecting a treatment preferably for at least 1 month, preferably for at least 3 months, preferably for at least 6 months,  
20 preferably for at least 9 months, and preferably for at least 12 months.

Acceptable protocols to administer therapeutic compositions in an effective manner include individual dose size, number of doses, frequency of dose administration, and mode of administration. Determination of such protocols can be accomplished by those skilled in the art. A suitable single dose is a dose that is capable of protecting an  
25 animal from disease when administered one or more times over a suitable time period. For example, a preferred single dose of a protein, mimetope or antibody therapeutic composition, including a recombinant protein vaccine, is from about 1 microgram ( $\mu\text{g}$ ) to

about 10 milligrams (mg) of the therapeutic composition per kilogram body weight of the animal. Booster vaccinations can be administered from about 2 weeks to several years after the original administration. Booster administrations preferably are administered when the immune response of the animal becomes insufficient to protect the animal from disease. A preferred administration schedule is one in which from about 10  $\mu$ g to about 1 mg of the therapeutic composition per kg body weight of the animal is administered from about one to about two times over a time period of from about 2 weeks to about 12 months. Modes of administration can include, but are not limited to, subcutaneous, intradermal, intravenous, intranasal, oral, transdermal, intraocular, intranasal, conjunctival, and intramuscular routes. Methods of administration for other therapeutic compounds can be determined by one skilled in the art, and may include administration of a therapeutic composition one or more times, on a daily, weekly, monthly or yearly regimen; routes of administration can be determined by one skilled in the art, and may include any route. A preferred route of administration of an inhibitory compound when administering to fleas is a topical, or "spot-on" formulation administered to the body surface of the animal, so that a flea would encounter the inhibitory compound when attached to the animal; another preferred route of administration of an inhibitory compound is an oral formulation that, when fed to an animal, would enter the bloodstream of the animal, which would then be transferred to a flea while feeding from the animal.

A recombinant protein vaccine of the present invention comprises a recombinantly-produced flea HMT and/or HNC protein of the present invention that is administered to an animal according to a protocol that results in the animal producing a sufficient immune response to protect itself from a flea infestation. Such protocols can be determined by those skilled in the art.

According to one embodiment, a nucleic acid molecule of the present invention can be administered to an animal in a fashion to enable expression of that nucleic acid

molecule into a protective protein or protective RNA (e.g., antisense RNA, ribozyme, triple helix forms or RNA drug) in the animal. Nucleic acid molecules can be delivered to an animal in a variety of methods including, but not limited to, (a) administering a naked (i.e., not packaged in a viral coat or cellular membrane) nucleic acid as a genetic  
5 vaccine (e.g., as naked DNA or RNA molecules, such as is taught, for example in Wolff et al., 1990, *Science* 247, 1465-1468) or (b) administering a nucleic acid molecule packaged as a recombinant virus vaccine or as a recombinant cell vaccine (i.e., the nucleic acid molecule is delivered by a viral or cellular vehicle).

A genetic (i.e., naked nucleic acid) vaccine of the present invention includes a  
10 nucleic acid molecule of the present invention and preferably includes a recombinant molecule of the present invention that preferably is replication, or otherwise amplification, competent. A genetic vaccine of the present invention can comprise one or more nucleic acid molecules of the present invention in the form of, for example, a dicistronic recombinant molecule. Preferred genetic vaccines include at least a portion of  
15 a viral genome, i.e., a viral vector. Preferred viral vectors include those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, picornaviruses, and retroviruses, with those based on alphaviruses, such as sindbis or Semliki forest virus, species-specific herpesviruses and poxviruses being particularly preferred. Any suitable transcription control sequence can be used, including those disclosed as suitable for protein  
20 production. Particularly preferred transcription control sequences include cytomegalovirus immediate early (preferably in conjunction with Intron-A), Rous sarcoma virus long terminal repeat, and tissue-specific transcription control sequences, as well as transcription control sequences endogenous to viral vectors if viral vectors are used. The incorporation of a "strong" polyadenylation signal is also preferred.

25 Genetic vaccines of the present invention can be administered in a variety of ways, with intramuscular, subcutaneous, intradermal, transdermal, conjunctival, intraocular, intranasal and oral routes of administration being preferred. A preferred

single dose of a genetic vaccine ranges from about 1 nanogram (ng) to about 600  $\mu$ g, depending on the route of administration and/or method of delivery, as can be determined by those skilled in the art. Suitable delivery methods include, for example, by injection, as drops, aerosolized and/or topically. Genetic vaccines of the present invention can be  
5 contained in an aqueous excipient (e.g., phosphate buffered saline) alone or in a carrier (e.g., lipid-based vehicles).

A recombinant virus vaccine of the present invention includes a recombinant molecule of the present invention that is packaged in a viral coat and that can be expressed in an animal after administration. Preferably, the recombinant molecule is  
10 packaging- or replication-deficient and/or encodes an attenuated virus. A number of recombinant viruses can be used, including, but not limited to, those based on alphaviruses, poxviruses, adenoviruses, herpesviruses, picornaviruses, and retroviruses. Preferred recombinant virus vaccines are those based on alphaviruses (such as Sindbis virus), raccoon poxviruses, species-specific herpesviruses and species-specific  
15 poxviruses. An example of methods to produce and use alphavirus recombinant virus vaccines are disclosed in U.S. Patent No. 5,766,602 to Xiong and Grieve, which is incorporated by reference herein in its entirety.

When administered to an animal, a recombinant virus vaccine of the present invention infects cells within the immunized animal and directs the production of a  
20 protective protein or RNA nucleic acid molecule that is capable of protecting the animal from flea infestation as disclosed herein. For example, a recombinant virus vaccine comprising a flea HMT and/or HNC nucleic acid molecule of the present invention is administered according to a protocol that results in the animal producing a sufficient immune response to protect itself from flea infestation. A preferred single dose of a  
25 recombinant virus vaccine of the present invention is from about  $1 \times 10^4$  to about  $1 \times 10^8$  virus plaque forming units (pfu) per kilogram body weight of the animal. Administration protocols are similar to those described herein for protein-based vaccines, with



subcutaneous, intramuscular, intranasal, intraocular, conjunctival, and oral administration routes being preferred.

A recombinant cell vaccine of the present invention includes recombinant cells of the present invention that express at least one protein of the present invention. Preferred  
5 recombinant cells for this embodiment include *Salmonella*, *E. coli*, *Listeria*, *Mycobacterium*, *S. frugiperda*, yeast, (including *Saccharomyces cerevisiae* and *Pichia pastoris*), BHK, CV-1, myoblast G8, COS (e.g., COS-7), Vero, MDCK and CRFK recombinant cells. Recombinant cell vaccines of the present invention can be administered in a variety of ways but have the advantage that they can be administered  
10 orally, preferably at doses ranging from about  $10^8$  to about  $10^{12}$  cells per kilogram body weight. Administration protocols are similar to those described herein for protein-based vaccines. Recombinant cell vaccines can comprise whole cells, cells stripped of cell walls or cell lysates.

The efficacy of a therapeutic composition of the present invention to protect an  
15 animal from flea infestation can be tested in a variety of ways including, but not limited to, detection of protective antibodies (using, for example, proteins or mimetopes of the present invention), detection of cellular immunity within the treated animal, or challenge of the treated animal with the fleas to determine whether the treated animal is resistant to infestation. Challenge studies can include direct administration of fleas to the treated  
20 animal. In one embodiment, therapeutic compositions can be tested in animal models such as mice. Such techniques are known to those skilled in the art.

One therapeutic composition of the present invention includes an inhibitor of flea HMT and/or HNC protein activity, i.e., a compound capable of substantially interfering with the function of a flea HMT and/or HNC protein susceptible to inhibition by an  
25 inhibitor of flea HMT and/or HNC protein activity. An inhibitor of HMT and/or HNC protein activity can be identified using flea HMT and/or HNC proteins of the present invention. An inhibitor of HMT and/or HNC protein function can be identified using flea

HMT and/or HNC proteins of the present invention. A preferred inhibitor of HMT and/or HNC protein function is a compound capable of substantially interfering with the function of a flea HMT and/or HNC protein and which does not substantially interfere with host animal proteins. As used herein, a compound that does not substantially inhibit or interfere with host animal proteins is one that, when administered to a host animal, the host animal shows no significant adverse effects attributable to the compound and which, when administered to an animal in an effective manner, is capable of protecting that animal from flea infestation.

One embodiment of the present invention is a method to identify a compound capable of inhibiting HMT and/or HNC protein activity of a flea. Such a method includes the steps of (a) contacting (e.g., combining, mixing) an isolated flea HMT and/or HNC protein, preferably a *C. felis* HMT and/or HNC protein of the present invention, with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has HMT and/or HNC protein activity, and (b) determining if the putative inhibitory compound inhibits the activity. HMT and/or HNC protein activity can be determined in a variety of ways known in the art, including but not limited to determining the ability of HMT and/or HNC protein to bind to or otherwise interact with a substrate. Such conditions under which a HMT and/or HNC protein has HMT and/or HNC protein activity include conditions in which a HMT and/or HNC protein has a correct three-dimensionally folded structure under physiologic conditions, i.e. physiologic pH, physiologic ionic concentrations, and physiologic temperatures.

Putative inhibitory compounds to screen include antibodies (including fragments and mimetopes thereof), putative substrate analogs, and other, preferably small, organic or inorganic molecules. Methods to determine HMT and/or HNC protein activity are known to those skilled in the art; see, for example, the Examples section of the present application. Methods to determine binding of a putative inhibitory compound to a HMT and/or HNC protein of the present invention are known to those of skill in the art and

include, for example, determining changes in molecular mass using surface plasmon resonance (e.g., determining light scatter by an inhibitor of a HMT and/or HNC protein, before and after contacting the inhibitor or protein with a HMT and/or HNC protein or inhibitor, respectively) or screening for compounds that inhibit interaction between a  
5 HMT and/or HNC protein and a substrate.

A preferred method to identify a compound capable of inhibiting HMT and/or HNC protein activity includes contacting an isolated flea HMT and/or HNC protein encoded by a nucleic acid molecule of Table I and/or Table II; (b) a protein comprising an at least 25 consecutive amino acid portion identical in sequence to a consecutive  
10 amino acid portion of a sequence as set forth in (a), wherein the protein has HMT and/or HNC protein activity; (c) a protein comprising a fragment of a protein as set forth in (a), wherein the fragment has an activity selected from the group consisting of binding to a HMT and/or HNC molecule and hydrolyzing a HMT and/or HNC protein substrate; and (d) a protein encoded by an allelic variant of a nucleic acid molecule that encodes any  
15 protein of (a), (b), or (c), with a putative inhibitory compound under conditions in which, in the absence of the compound, the protein has HMT and/or HNC protein activity; and determining if the putative inhibitory compound inhibits the activity.

Another embodiment of the present invention is an assay kit to identify an inhibitor of a flea HMT and/or HNC protein of the present invention. This kit comprises  
20 an isolated flea HMT and/or HNC protein of the present invention, and a means for determining inhibition of an activity of flea HMT and/or HNC protein, where the means enables detection of inhibition. Detection of inhibition of flea HMT and/or HNC protein identifies a putative inhibitor to be an inhibitor of flea HMT and/or HNC protein. Means for determining inhibition of flea HMT and/or HNC protein include an assay system that  
25 detects binding of a putative inhibitor to a flea HMT and/or HNC molecule, and an assay system that detects interference by a putative inhibitor of the ability of flea HMT and/or

HNC protein to hydrolyze a substrate. Means and methods are described herein and are known to those skilled in the art.

The following examples are provided for the purposes of illustration and are not intended to limit the scope of the present invention. The following examples include a number of recombinant DNA and protein chemistry techniques known to those skilled in the art; see, for example, Sambrook et al., *ibid*.

#### Example 1

This Example describes the isolation of RNA from the hindgut and Malpighian tubules (HMT) of *Ctenocephalides felis* and the use of isolated RNA to construct subtracted and unsubtracted cDNA libraries.

Approximately 10,000 hindguts and Malpighian tubules were dissected from equal numbers of cat blood fed and unfed adult *C. felis* with a male to female ratio of 1 to 4, and total RNA was extracted using a guanidine isothiocyanate lysis buffer and the standard procedure described by Sambrook et al. Poly-A enriched mRNA was purified from total RNA above using a mRNA Purification Kit, available from Pharmacia Biotech, Piscataway, NJ, following the manufacturer's protocol. The same procedures were used to extract total RNA and isolate poly-A enriched mRNA from the dissected *C. felis* bodies following removal of HMT, referred to hereinafter as "non-HMT mRNA".

Poly-A enriched mRNA was used to construct a cDNA library using subtractive hybridization and suppression PCR as follows. Subtractive hybridization and suppression PCR was conducted using a PCR-Select™ cDNA Subtraction Kit, available from Clontech Laboratories, Inc., Palo Alto, CA according to the manufacturer's instructions. Briefly, this kit uses subtractive hybridization and suppression PCR to specifically amplify cDNA sequences that are present in the tester cDNA and absent in the driver cDNA, thus enriching for tester-specific sequences. The efficiency of the subtraction process can be assessed by semi-quantitative PCR and by comparing the ethidium bromide staining patterns of the subtracted and unsubtracted samples on agarose

gels as described in section V.D. of the manufacturer's protocol. For the semi-quantitative PCR, three genes with mRNAs known to be expressed outside of the HMT tissue were used to test for specific subtraction. These genes encoded putative actin, N-aminopeptidase, and serine protease proteins.

5           Subtractive hybridization and suppression PCR was conducted under the following conditions. Two micrograms ( $\mu\text{g}$ ) of HMT mRNA was used as the template for synthesis of the tester material and 2  $\mu\text{g}$  of non-HMT mRNA was used as template for synthesis of the driver material in this reaction. The number of cycles used in the selective amplification steps was optimized using the manufacturer's protocols.

10          Optimization resulted in the use of 24 rather than the standard 27 cycles of primary PCR in combination with 15 cycles of secondary PCR rather than the standard 12 cycles.

            The products from the suppressive PCR reaction were ligated into the pCR<sup>®</sup>2.1 vector, available from Invitrogen, Carlsbad, CA, using an Original TA Cloning<sup>®</sup> Kit, available from Invitrogen. The ligation reaction was then used to transform INV $\alpha$ F' One

15       Shot<sup>™</sup> competent cells, available from Invitrogen, which were plated on Luria broth (LB) agar with 50 micrograms per milliliter ( $\mu\text{g}/\text{ml}$ ) ampicillin, available from Sigma-Aldrich Co., St. Louis, MO, and 50  $\mu\text{g}/\text{ml}$  5-bromo-4-chloro-3-indoyl  $\beta$ -D-galactopyranoside (X-Gal), available from Fisher Biotech, Fair Lawn, NJ. Transformed colonies were amplified and the DNA isolated using the standard alkaline lysis procedure described by

20       Sambrook et al., *ibid*.

            Automated cycle sequencing of DNA samples was performed using an ABI PRISM<sup>™</sup> Model 377, available from Perkins Elmer, with XL upgrade DNA Sequencer, available from PE Applied Biosystems, Foster City, CA, after reactions were carried out using the PRISM<sup>™</sup> Dye Terminator Cycle Sequencing Ready Reaction Kit or the PRISM<sup>™</sup>

25       dRhodamine Terminator Cycle Sequencing Ready Reaction Kit or the PRISM<sup>™</sup> BigDye<sup>™</sup> Terminator Cycle sequencing Ready Reaction Kit, available from PE Applied Biosystems, following the manufacturer's protocol, hereinafter "standard sequencing

methods". The resulting sequences are presented in Table II. Sequence analysis was performed using the MacVector<sup>™</sup> sequence analysis software, available from International Biotechnologies Inc., New Haven, CT, and the SeqLab sequence analysis software, using default parameters. Each sequence read was trimmed of vector sequence at either end and submitted for a search through the National Center for Biotechnology Information (NCBI), National Library of Medicine, National Institute of Health, Baltimore, MD, using the BLAST network. This database includes SwissProt + PIR + SPupdate + GenPept + GUpdate + PDB databases. The search was conducted using the xBLAST function, which compares the translated sequences in all 6 reading frames to the protein sequences contained in the database.

An unsubtracted HMT cDNA library was constructed as follows. Approximately 10,000 HMT tissues were dissected from equal numbers of unfed and cat blood-fed adult *C. felis* with a male to female ratio of 1:4. Total RNA was extracted using a guanidine isothiocyanate lysis buffer and procedures described in Sambrook et al., followed by isolation using a mRNA purification kit, available from Pharmacia, according to the manufacturer's protocols. The library was constructed with 5 µg of isolated mRNA using a ZAP-cDNA<sup>®</sup> cDNA synthesis kit, and packaged using a ZAP-cDNA<sup>®</sup> Gigapack<sup>®</sup> gold cloning kit, both available from Stratagene, La Jolla, CA. The resultant HMT library was amplified to a titer of about  $5 \times 10^9$  plaque forming units per milliliter (pfu/ml). Single clone excisions were performed using the Ex-Assist<sup>™</sup> helper phage, available from Stratagene, and used to create double stranded plasmid template for sequencing using the manufacturer's protocols with the following exceptions. Following incubation of the SOLR cells with the cleared phage lysate, the mixture was used to inoculate LB broth, and the mix was incubated overnight and then subjected to mini-prep plasmid preparation and sequencing as described for the subtracted HMT library above.

## Example 2

This Example describes the isolation of RNA from the head and nerve cord (HNC) of *Ctenocephalides felis* and the use of isolated RNA to construct subtracted and unsubtracted cDNA libraries.

5            Approximately 4,000 heads and attached nerve cords, including the terminal abdominal ganglia were dissected from equal numbers of cat blood-fed and unfed adult *C. felis* with a male to female ratio of 1 to 4, and total RNA was extracted using a guanidine isothiocyanate lysis buffer and the standard procedure described by Sambrook et al. Approximately 618 µg of total RNA was recovered. Poly-A enriched mRNA was  
10 purified from total RNA above using a mRNA Purification Kit, available from Pharmacia, following the manufacturer's protocol. Approximately 13 µg of mRNA was isolated. The same procedures were used to extract total RNA and isolate poly-A enriched mRNA from the dissected *C. felis* bodies following removal of HNC tissues, referred to hereinafter as "non-HNC mRNA".

15            Suppression subtractive PCR was conducted as described in Example 1 using a PCR-Select™ cDNA Subtraction kit, available from Clontech, under the following conditions. Two micrograms (µg) of HNC mRNA was used as the template for synthesis of the tester material and 2 µg of non-HMT mRNA was used as template for synthesis of the driver material in this reaction. The number of cycles used in the selective  
20 amplification steps was optimized using the manufacturer's protocols. Optimization resulted in the use of 24 rather than the standard 27 cycles of primary PCR in combination with either 12 or 15 cycles of secondary PCR.

cDNA pools from various PCR cycling combinations were ligated into the TA vector using a TA cloning kit, available from Invitrogen. Aliquots of ligation reaction  
25 were transformed into Ultramax DH5α™ bacteria, available from Gibco-BRL, Gaithersburg, MD. Portions of the transformation mixes were used to inoculate LB broth cultures containing 100 µg/ml of ampicillin. The overnight cultures were plated to

generate discreet colonies which were used individually for overnight cultures for plasmid preps. Transformed colonies were amplified and the DNA isolated using the standard alkaline lysis procedure described by Sambrook et al., *ibid*.

Automated cycle sequencing of DNA samples was performed using the standard  
5 sequencing methods described in Example 1. The resulting sequences are presented in Table I. Sequence analysis was performed using the MacVector™ sequence analysis software, available from International Biotechnologies Inc., New Haven, CT, and the SeqLab sequence analysis software, using default parameters.

An unsubtracted cDNA library was constructed as follows. Approximately 6400  
10 head and nerve cords were dissected from *C. felis* and poly-A RNA was isolated as described above. About seven µg of HNC poly-A RNA was used to construct a cDNA library using Stratagene's λZAP-cDNA Synthesis Kit and protocol. The resultant HNC library was amplified to a titer of about  $5 \times 10^9$  plaque forming units per milliliter (pfu/ml). Single clone excisions were performed using the Ex-Assist helper phage,  
15 available from Stratagene, and used to create double stranded plasmid template for sequencing using the manufacturer's protocols with the following exceptions. Following incubation of the SOLR cells with the cleared phage lysate, the mixture was used to inoculate LB broth, and the mix was incubated overnight and then subjected to mini-prep plasmid preparation and sequencing as described for the subtracted library above.

20 While various embodiments of the present invention have been described in detail, it is apparent that modifications and adaptations of those embodiments will occur to those skilled in the art. It is to be expressly understood, however, that such modifications and adaptations are within the scope of the present invention, as set forth in the following claims: